

Asian Journal of Food Research and Nutrition

Volume 3, Issue 4, Page 1203-1213, 2024; Article no.AJFRN.127515

Carcass, Internal Organ Characteristics and GIT Morphometry of Broiler Chickens Fed Diets Containing Fermented Sweet Orange (*Citrus sinensis*) Peel Meal (FSOPM) as a Replacement of Maize

Akpe, M.E. ^{a*}, Oluremi, O.I.A. ^b and Tuleun, C.D. ^c

^a Department of Food Science and Technology, University of Mkar, Mkar, Nigeria. ^b Department of Animal Science, The National University of Lesotho, Roma, Lesotho. ^c Department of Animal Nutrition, Joseph Sarwuan Tarka University, Makurdi, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/ajfrn/2024/v3i4210

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/127515

> Received: 25/09/2024 Accepted: 29/11/2024 Published: 04/12/2024

Original Research Article

ABSTRACT

The experiment was carried out to determine the effect of fermented sweet orange peel meal (FSOPM) as a replacement of maize on carcass, internal organs characteristics and GIT morphometry of broiler chicken. Fresh sweet orange peel (SOP), and fresh rumen content (RC)

*Corresponding author: E-mail: msurshimaakpe@gmail.com;

Cite as: M.E., Akpe, Oluremi, O.I.A., and Tuleun, C.D. 2024. "Carcass, Internal Organ Characteristics and GIT Morphometry of Broiler Chickens Fed Diets Containing Fermented Sweet Orange (Citrus Sinensis) Peel Meal (FSOPM) As a Replacement of Maize". Asian Journal of Food Research and Nutrition 3 (4):1203-13. https://doi.org/10.9734/ajfrn/2024/v3i4210.

were collected and thoroughly mixed to obtain a homogenous mixture in a ratio of 1: 0.3 (SOP:RC) respectively. The mixture was fermented for 48 hours and sun-dried to < 10% moisture. The fermented SOP was milled to obtain FSOPM. The FSOPM was sub-sampled and its proximate composition determined, after which it was included at 0%, 2%, 4%, 6%, 8%, and 10% respectively in the diets. 216 day – old Marshal MY broiler chicks were randomly assigned to 6 dietary treatments in a complete randomized design (CRD). Each treatment was replicated 3 times with 2 birds in each replicate. The birds were fed the test diets for 56 days and carcass, internal organs and GIT were evaluated. The fasted weight, eviscerated weight, dressed weight, dressing percentage, thigh, wing, head, shank, abdominal fat, empty gizzard, liver, kidney, proventriculus, bile, GIT weight, large intestine and small intestine were significantly different (p < 0.05). Inclusion of FSOM increased carcass cut, internal organs and the GIT morphometry; and recessed abdominal fat pad of broiler chickens. It was concluded that 10% FSOPM replacement of maize was safe and optimum for broiler chicken production.

Keywords: Fermentation; sweet orange peels; rumen content; Broiler chickens; proximate analysis; growth performance.

1. INTRODUCTION

Agro-industrial waste products are in abundance in Nigeria and industries producing these byproducts often incure expenses for their proper disposal. The processing of orange fruits (citrus) generates significant amounts of by-products often containing valuable compounds in the peels, pulp and seeds. In 2018, the United Nations Food and Agriculture Organization [1] estimated a world citrus production of 104.15Mt, with 75.54Mt corresponding to orange. The waste obtained from industrial processed citrus is made up of valuable by-products which according to [2] contains 89.20% DM, 8.20% CP, 13.30% CF, 4.51% EE, 6.09% ash, 67.90% NFE and 3079.61 Kcal/kg of ME that can be used for poultry feed. Harnessing these wastes will reduce the pressure of demand, as well as cost of maize, the major and most widely utilized energy ingredient in poultry diets which have risen over the years [3 and 4]. The high cost of animal feeding results from the various use to which maize, a conventional feed stuff is being competed for as a staple food for man, in brewing and confectionary. According to [5 and 6] the availability of high-quality poultry feeds is limited by phenomenal rise in the cost of conventional feed ingredient for animals, mostly due to competition from direct consumption of these materials by man. For a feed stuff to be qualified and used as an alternative for livestock feeding, the ingredient most be cheap, readily available and not used as a staple food for humans, but should meet the nutritional requirements of farm animals [7, 8 and 9]. Limitations of the use of sweet orange peel

(SOP) has been reported due to some antinutritional factors such as tannin, saponin, phytate, oxalate, flavoniod and limonene [10]. It is important therefore that certain processing techniques be employed to minimize these antinutritional factors and improve the nutritive value of SOP.

The rumen content (RC), an abattoir waste is the material obtained from the rumen of slaughtered ruminants. It contains a mixed population of microbes and undigested feed material and can be used as a natural inoculum. Fermentation is a unique process with great potentials of recycling agro by-products such as sweet orange peels into useful poultry feeds in developing countries. The process is easy to manage on farm conditions and does not require the use of chemicals. Fermentation eliminates toxins, improves nutritional quality. increases digestibility, palatability, bioavailability of minerals and simple sugars [10 and 11]. This experiment was therefore carried out to determine the effect of fermented sweet orange peel meal (FSOPM) on carcass yield, internal organs characteristics and GIT morphometry of broiler chickens.

2. MATERIALS AND METHODS

2.1 Experimental Site

The experiment was conducted at the poultry section of the Livestock Unit, on the Teaching and Research Farm, Joseph Sarwuan Tarka University, Makurdi Benue State, Nigeria. The area is warm with a minimum temperature range of $24.20 \pm 1.40^{\circ}$ C and a maximum temperature range of $36.33 \pm 3.70^{\circ}$ C [12].

		Treatments				
Ingredients	0%	2%	4%	6%	8%	10%
-	FSOPM	FSOPM	FSOPM	FSOPM	FSOPM	FSOPM
Maize	56.3	55.17	54.05	52.92	50.80	50.67
FSOPM	0.00	1.13	2.25	3.38	4.50	5.63
SBM	32.45	32.45	32.45	32.45	32.45	32.45
BDG	5.00	5.00	5.00	5.00	5.00	5.00
Bone ash	2.20	2.20	2.20	2.20	2.20	2.20
Limestone	1.50	1.50	1.50	1.50	1.50	1.50
Blood meal	1.50	1.50	1.50	1.50	1.50	1.50
Methionine	0.25	0.25	0.25	0.25	0.25	0.25
Lysine	0.30	0.30	0.30	0.30	0.30	0.30
Common salt	0.25	0.25	0.25	0.25	0.25	0.25
Vt/min pmx	0.25	0.25	0.25	0.25	0.25	0.25
Total	100.00	100.00	100.00	100.00	100.00	100.00
Calculated	Analysis					
Crude protein (%)	21.77	26.76	21.82	21.73	21.62	21.70
Crude fibre (%)	4.23	4.35	4.49	4.60	4.68	4.82
Fat (%)	3.76	3.75	3.78	3.74	3.68	3.82
ME (Kcal/kg)	2065.86	2385.98	2865.09	3019.62	2019.61	2050.94
Methionine (%)	0.61	0.60	0.61	0.60	0.60	0.59
Lysine (%)	1.45	1.44	1.44	1.44	2.43	1.43
Calcium (%)	1.44	1.44	1.44	1.44	1.43	1.43
Phosphorus (%)	0.62	0.63	0.65	0.62	0.61	0.61

Table 1. Composition of Broiler Finisher Diets Containing FSOPM

Vitamin – mineral premix* (animal care ®) supplied the following nutrients per kg of feed. Vitamin: A (1,200,000 I.U) D₃ (300,000 I.U), E (3,000mg), K₃ (250mg), folic acid (10mg), Niacin (4,000mg), Calpan (1,000mg), B₂ (500mg), B₁₂ (2mg), B1 (200mg), B₆ (350mg), Biotin (8mg), Antioxidant (12,500mg), Minerals; Cobalt (25mg), Selenium (25mg), Iodine (120mg), Iron (4,000mg), Manganese (700mg), Copper (800mg), Zinc (6,000mg), and Chlorine chloride (20,000mg) FSOPM=fermented sweet orange peel meal, SBM= soybean meal, BDG=brewer's dried grains, ME=metabolizable energy, vit/min pmx= vitamins and mineral premixes

2.2 Preparation of Fermented Sweet Orange Peel, Rumen Content and Diets

Sweet orange peels were collected together as a composite irrespective of variety from sweet orange fruit retailers while the rumen content (RC) was collected from freshly slaughtered white Fulani cattle at the slaughter slab all in Gboko, Benue State. The Sweet Orange Peels (SOP) were thoroughly mixed with the rumen content (RC) to obtain a homogenous mixture in a ratio of 1: 0.3. The mixture was packed, tied in polythene bags and allowed to ferment for 48 hours. The fermented materials were sun-dried, separated from the RC and the fermented peels were milled to obtain fermented sweet orange peel meal (FSOPM). The fermented Sweet Orange peel meal was then sub-sampled and the composition determined following proximate standard procedure [13]. Phytate was determined using spectrophotomeric method [14] tannin and saponin using the procedure of [15], Oxalate using the technique of [16]. The FSOPM was then incorporated in broiler diets for starter and finisher phases at 0, 2, 4, 6, 8 and 10% to produce 0% FSOPM (control), 2% FSOPM, 4% FSOPM, 6% FSOPM, 8% FSOPM and 10% FSOPM diets presented in Tables 2 and 3 respectively.

2.3 Experimental Broilers and Design

Two hundred and sixteen (216) day – old Marshall MY broiler chicks were purchased from Chi Farms, Ibadan, balanced for weight and randomly assigned to six dietary treatments in a complete randomized design (CRD). Each treatment group of 36 broilers was further subdivided into three replicates of 12 chicks each. The broiler birds were reared in deep litter, halfwalled open – sided house with the upper half covered with wire mesh for good ventilation. Feed and water were provided *ad libitum* throughout the feeding trial which lasted for 56 days. Standard management procedures were observed. Feed intake and body weights were recorded weekly. Feed intake was determined by serving birds weighed amount of feed and the corresponding left-over obtained by difference and daily feed intake calculated as average for seven days.

2.4 Data Collection and Analysis

a). Carcass evaluation

Carcass evaluation was carried out using the procedure of [17]. At the termination of the feeding trial on the 56th day, one finisher broiler per replicate whose live weight similar to the group's average weight was selected, fasted for 18 hours, slaughtered by severing the head from the neck with a sharp knife and bled. The broilers were dipped in hot water ($55 - 60^{\circ}$ C) defeathered and carcass separated into cuts namely breast, drumstick, thigh, back, wing and neck. Offal's separated included head, shank and abdominal fat. Carcass weight was determined by weighing individual cuts, and each expressed as percent of live weight (% LW).

b). Internal organs

Each of the broilers used in carcass evaluation, had its internal organs examined as well. Internal organs namely; empty gizzard, liver, lungs, kidney, proventriculus, pancreases, bile and spleen were separated and weighed using an electronic scale. Gastro-intestinal tracts of slaughtered birds were evaluated by taking the weights of the GIT, the small intestine, the large intestine and the caecum. The length of the GIT and its components (the small intestine, the large intestine and the caecum) were also measure.

2.5 Data Analysis

Data generated were subjected to one-way analysis of variance (ANOVA) using Minitab software [18], and where significant differences occurred, means were separated using the least significant difference (LSD) as contained in [18].

3. RESULTS AND DISCUSSION

3.1 Proximate Composition

The proximate composition of fermented sweet orange peel meal (FSOPM) is presented in Table 2. For a feed stuff to be qualified for use as an alternative for poultry feeding, the nutrient value of such a material should be comparable nutritionally to that of the conventional feed stuff it replaces like maize and should also meet the nutritional requirements of broiler chickens.

Table 2. Proximate analysis, Energy and Antinutritional Factors in Fermented Sweet Orange Peel Meal (FSOPM)

Parameters (%)	(FSOPM)
Dry matter	91.92
Crude protein	7.82
Crude fibre	13.88
Ether extract	2.94
Ash	8.16
Nitrogen free extract	67.20
Metabolizable Energy (kcal/kg)	2915.42
Tannin	0.008
Saponin	0.70
Alkaloid	2.50
Phytate	0.45
Oxalate	3.02
	, ,

FSOPM=fermented sweet orange peel meal

The dry matter (DM) value (91.92%) obtained in this study is comparable to 92.79% - 93.22% reported by [19] for sun-dried SOP, but higher than 81.43% - 87.05% and 89.61 - 92. 56% reported by [11] and [20] respectively. The DM content was above 90% indicating that all the nutrients were seemingly high. This high DM content will enhance long shelf - life storage of FSOPM. [21] reported that high DM favoured long storage duration of BSOPM. The crude protein value (7.82%) obtained in this study is comparable to 7.71% - 9.26% reported by [11] but higher than 7.22%, 6.44% and 6.78% -7.30% reported by [22, 23 and 19] respectively. The crude protein value in FSOPM was not far from that of maize (8.90%), a conventional energy feed stuff [24] suggestive that, it can partially be used in place of maize. Crude fibre value (13.88%) obtained in this study compared favourably with 14.73%, 13.30% and 15.77% reported by [11] respectively. The CF in FSOPM was higher than that of maize (2.70%) as reported by [24], but lower than 14.60% reported by [11] for sweet orange peels (SOP). The reduced CF percent in this study could be attributed to fermentation. Maize been a parent /conventional feed material, it was also expected that the CF of FSOPM as a by-product, be higher. The ether extract value of FSOPM (2.94%) obtained in this study compares favorably to 2.23% - 3.41% and 2.50% - 2.95% reported by [19], when sweet orange fruit peels were fermented. The values were however higher than 1.88% - 2.65% and 1.96% reported by [22 and 25] respectively. The ash value of FSOPM (8.16%) obtained in this study was comparable to 8.19% [19] and 9.91% [2] but higher than 6.09% reported by [2]. The ash content of FSOPM was higher than 2.70% for maize [24]. Increase in ash content could be attributed to the rumen content used for fermentation. The result shows that FSOPM contains more total minerals than maize (a conventional energy feedstuff) and could perfectly replace maize on the grounds of mineral supply. [26] had earlier reported that grains are relatively poor in minerals. Nitrogen free extract (NFE) which represents the soluble carbohydrates (67.20%) obtained in this study compared favourably to 67.90% reported by [2]. The value for NFE obtained in this study was lower than 83.00% [27] but higher than 50.75% reported by [23] for maize. This result suggested that maize contains more utilizable sugars than FSOPM. Typical of agricultural by-products one of their nutritional characteristics is the low energy/NFE content. In comparism to maize, his result was also expected. The ME value of FSOPM (2915.42 Kcal/kg) obtained in this study was comparable to 2795.28 - 2913.92 Kcal/kg reported by [11] but was lower than 3752.12 Kcal/kg reported by [27] for maize, and 3432.32 Kcal/kg reported for maize [28]. The ME obtained though inferior to maize can be used as a source of energy in the diet of broiler chicken. [29] had earlier reported that SOP could serve mainly as an energy source because of its high energy value (2990.00 Kcal/kg). The energy value of FSOPM in this study suggests its suitability and utilization as an energy source in broiler diets

3.2 Carcass Cuts and Offals

The results of carcass cuts (breast, drumstick, thigh, back, wing, neck) and offals (head, shank abdominal fat) are presented in Table 3. There was no significant difference (p> 0.05) in the breast, drumstick, back and neck. The thigh cut was significantly higher (p< 0.05) at 0% and 10% FSOPM and significantly lower (p< 0.05) at 4% FSOPM. The wing was significantly higher (p < p0.05) at 10% FSOPM and significantly lower (p < p0.05) at 6% FSOPM. The live weight (1.59 - 2.00 kg) obtained in this study was comparable to 1.33 - 2.01 kg reported by [30], and the average of 1.92 kg reported by [31]. The average live weight obtained in this study was within broiler chicken Table 3 weight range of 1600 g to 2000 g reported by [32]. The dressing percent (62.09% - 69.00%) obtained in this study was comparable to 72.41% reported by [22] but was higher than 62.29% reported by [31]. The dressing percent

obtained in this study agrees with 71.00% - 75% earlier reported by [33] as satisfactory for broiler. Dressing percentage is the absolute value of saleable meat which therefore showed that the chickens in the FSOPM dietary treatment groups produced lean meat for utilization. The thigh weight (14.70% - 16.65%) obtained in this study was comparable to 15.34% - 17.55% reported by [34] when broiler chickens were fed rice milling waste (RMW) based diets. The thigh weight was however higher than 11.55% - 13.15% reported by [22] when broiler chickens were fed water soaked sweet orange fruit peel-based diets. The wing weight (12.16% - 13.62%) obtained in this study was comparable to 10.77% - 14.35% reported by [34] but was higher than 7.81% -8.14% reported by [22] for broiler chickens. The thigh, breast and drumstick cuts are prime cuts of chicken and they give a picture of the carcass meatiness and eventual revenue vield. Carcass quality is closely related to the intake level of nutrients especially protein and energy. The values obtained in this study for carcass cuts of broiler chickens fed FSOPM are statistically similar to the maize based control diet. Implying therefore that, birds fed diets containing FSOPM yielded meat at levels equivalent to those on the control. The head and shank were significantly higher (p < 0.05) at 10% FSOPM; while abdominal fat was significantly higher (p < 0.05) at 0% FSOPM (control diet) and significantly lower (p > 0.05) at 6% and 10% FSOPM. The head weight (3.43% - 4.28%) and shank weight (5.74% - 7.34%) obtained in this study were comparable to (2.78% - 4.36%) but higher than (4.18% - 4.75%) for shank weight reported by [23] for sun-dried SOP supplemented with guantum blue enzyme. The abdominal fat (0.00% - 0.84%) obtained in this study was higher than 0.00 - 0.21% reported by [2] but lower than 1.85% - 2.13% reported by [35]. The result on abdominal fat clearly showed that the use of FSOPM in the diet of broiler chicken recessed the deposition of abdominal fat. This could be attributed to high fiber content of the diets and also to the various active components present in Citrus sinensis peels. According to [36] dietary addition of Citrus sinensis waste had beneficial effects in reducing the levels of cholesterol and fats in meat. This has good health benefits because fatty meat consumption can cause/or increase coronary challenges. This result is in agreement with those of [37] and [38] who reported significantly decrease in abdominal fat in broiler chicken fed diets supplemented with orange peel powder and biodegraded sweet orange peels, respectively.

Weight of part/	Treatments						
parameter (%)	0%	2%	4%	6%	8%	10%	_
	FSOPM	FSOPM	FSOPM	FSOPM	FSOPM	FSOPM	
Live weight (kg)	2.00 ^a	1.83 ^{abc}	1.77 ^{abc}	1.95 ^{ab}	1.73 ^{bc}	1.59°	0.08
Eviscerated weight	76.13ª	69.30°	72.41 ^b	75.24 ^{ab}	74.68 ^{ab}	73.54 ^{ab}	0.37
(%LW)							
Dressed weight (kg)	1.38ª	1.22 ^{ab}	1.21 ^{ab}	1.34 ^{ab}	1.17 ^{bc}	0.99 ^c	0.06
Dressed percent (%)	69.00 ^a	62.09 ^b	67.58ª	68.49 ^a	68.37ª	65.54ª	0.91
Breast (% LW)	29.93ª	29.03ª	30.34ª	30.38ª	30.08ª	28.03ª	1.00
Drumstick (% LW)	16.01ª	16.92ª	15.45ª	15.92ª	16.40 ^a	16.43ª	0.61
Thigh (% LW)	16.18ª	16.04 ^{ab}	14.70 ^b	15.98 ^{ab}	16.01 ^{ab}	16.65ª	0.55
Back (% LW)	14.83 ^a	15.52ª	15.92ª	15.18ª	14.65 ^a	14.92 ^a	0.57
Wing (% LW)	12.48 ^{bc}	12.97 ^{abc}	13.27 ^{ab}	12.16°	12.89 ^{abc}	13.62ª	0.30
Neck (% LW)	10.18ª	9.24 ^a	10.17ª	9.71ª	9.39 ^a	10.12ª	0.35
Head (% LW)	3.54 ^b	3.47 ^b	3.43 ^b	3.50 ^b	3.53 ^b	4.28 ^a	0.14
Shank (% LW)	6.72 ^{ab}	7.01 ^{ab}	5.74 ^c	6.34 ^{bc}	6.37 ^{bc}	7. 34 ^a	0.23
Abdominal fat (% LW)	0.84 ^a	0.57 ^{ab}	0.48 ^{ab}	0.06 ^b	0.43 ^{ab}	0.00 ^b	0.22

Table 3. Effect of FSOPM on Carcass parameters and Offals of Broiler Chickens

^{a,b,c,} Means with different superscripts in the same row are significantly different (p<0.05), FSOPM, fermented sweet orange peel meal, SEM, standard error of mean

3.3 Internal Organ Characteristics

Internal organ characteristics are presented in Table 4. Dietary treatment did not show any significant (p > 0.05) effect on internal organs like lungs, heart, pancreas and spleen. However, FSOPM had significant (p < 0.05) effect on empty gizzard, liver, kidney, proventriculus and bile. Variations in this study did not follow any definite pattern. The empty gizzard and liver were significantly higher (p < 0.05) at 10% FSOPM and significantly lower (p > 0.05) at 0% FSOPM. Broiler chickens on the FSOPM based diets had significantly (p < 0.05) heavier gizzard, proventriculus liver. kidnev and than chickens on the control diets; except for chickens on 6% FSOPM, which had lighter proventriculus.

The percent live weight of the empty gizzard (1.87% to 2.37%) obtained in this study is comparable to 2.17% reported by [39], but lower than the average of 2.81% reported by [2]. The significantly higher gizzard weight of the chickens on FSOPM based diets could be attributed to the higher CF content of the diets. The grinding process of these relatively more fibrous diets resulted to more muscular development of the gizzard thereby increasing its weight. This result is in agreement with the finding of [40] who reported that the inclusion of oat hulls in broiler diet at 5% stimulated the gizzard grinding activity, allowed better development of the muscular layers and caused an increase in the gizzard size. [41] stated that dietary fibre had

been seen to have a positive effect on gizzard development. The percent live weight of the proventriculus was higher in the test diets as FSOPM inclusion level increased from 2% to 6%. The values (0.44% to 0.65%) obtained were comparable to 0.59% and 0.49% reported by [31 and 39] respectively. It was however higher than 0.37% reported by [22]. The significant increase in the percent live weight of the proventiculus could be attributed to more digestive function the organ had to perform (increased secretion of pepsin and hydrochloric acid) in order to aid digestion of protein and also to initiate the breakdown of higher fiber content of the diet. [42] stated that diets high in fibre content may produce dilatation of the proventiculus with its increase in size and its contents. The percent live weight of the liver (1.88% - 2.53%) obtained in this study is comparable to 1.8% reported by [39], but higher than the value of 1.76% reported by [22]. The significant increase in the liver weight could be in positive response to increase secretion of the bile to aid emulsification of fats and thus facilitate the action of the enzyme lipase. High liver weight has also been shown to be an indication of higher metabolic rate of the liver to reduce or eliminate toxins in the body system [43]. As a major detoxifying organ, the increase in the size of the liver could be in attempt to detoxifying the residual anti-nutrients contained in FSOPM. The percent live weight of the kidney (0.56% - 0.89%) obtained in this study is comparable to 0.70% reported by [19], but was higher than 0.50% reported by [20]. The percent live weight of kidney decreased as the level of FSOPM in the diet increased from 2% to 6% and then increased at 8% inclusion level. [44] opined that anti-nutrients could cause increase kidney and liver weights owing to stress involved from detoxification. FSOPM contained anti-nutrients but all the values obtained in this study were within the safe recommended limits as reported by [45]. The percent live weight of the bile was higher for chickens on the FSOPM dietary treatments than the chickens on the maize based diet (control group) with the exception of FSOPM (2%), which appeared on outlier. The increase of the bile with increase in inclusion level of FSOPM in the diet could be that the liver secreted more bile to be stored and used for fat emulsification and hence the increase in size. The nonsignificant difference among the treatment groups for heart indicates normal blood circulation among the broilers in the dietary groups. Non -significant observations in the lungs, pancreas and spleen shows that birds tolerated the test ingredient and were not harmed by the different inclusion levels.

3.4 Gastro-intestinal Tract (GIT) Morphometry

Gastro-intestinal tract morphometry results are presented in Table 5. The GIT weight, large intestine weight and small intestine length were significantly affected (p < 0.05) among the treatment groups. Variation in this study did not follow any definite pattern. Other GIT components (weight or length) were not significantly affected (p > 0.05). The percent tract weight of the GIT (57.05 g - 72.96 g) was

comparable to 5.01% live weight reported by [19]. The broiler chickens on the FSOPM based diets had significantly (P < 0.05) heavier GIT weight than those on the control diet. The increase of the GIT weight could be attributed to increased energy expenditures. It has been previously reported that increase in the GIT size is intimately related with an increase of the energetic needs of the animal [46]. [42] stated that increase in the weight of the GIT is important to pancreatic enzymes secretion to achieve better growth in birds during the early stage of life.

The percent tract weight of the large intestine (6.24% - 13.63%) obtained was comparable to the average of 0.31% live weight reported by [19]. The small intestine (% GIT length) of broilers on FSOPM based diets was significantly (p < 0.050) longer than that of the control. This observation suggests a compensatory and inverse morphometric relationship between small intestine length and caecal length on one hand and the large intestine length on the other hand. This observation is in agreement with the findings of [47] who reported increase in the length of the small intestine and attributed it to the richness of soluble pectin contained in citrus pulp diets. This result also agrees with that of [48] who stated that supplementation of dried Citrus sinensis peel on the jejunum and ileum showed significant difference from that of the control group. The large intestine (% GIT weight) was significantly lighter (P < 0.05) at 2% FSOPM. Variation showed no definite pattern.

Parameter (% LW)	Treatments						
	0%	2%	4%	6%	8%	10%	
	FSOPM	FSOPM	FSOPM	FSOPM	FSOPM	FSOPM	
Live weight (kg)	2.00 ^a	1.83 ^{abc}	1.77 ^{abc}	1.95 ^{ab}	1.73 ^{bc}	1.59 ^{bc}	0.08
Empty gizzard	1.87 ^b	2.13 ^{ab}	2.21 ^{ab}	2.10 ^{ab}	2.36 ^a	2.37ª	0.15
Liver	1.88°	1.99 ^{bc}	2.11 ^{bc}	1.99 ^{bc}	2.27 ^{ab}	2.53ª	0.11
Lungs	0.53ª	0.58ª	0.66ª	0.66 ^a	0.54ª	0.64 ^a	0.06
Kidney	0.56 ^b	0.74 ^{ab}	0.74 ^{ab}	0.66 ^b	0.68 ^{ab}	0.89 ^a	0.06
Heart	0.46 ^a	0.49 ^a	0.45 ^a	0.42 ^a	0.46 ^a	0.50 ^a	0.03
Proventriculus	0.48 ^{ab}	0.53 ^{ab}	0.50 ^{ab}	0.44 ^b	0.48 ^{ab}	0.65ª	0.05
Pancreas	0.23ª	0.23ª	0.23 ^a	0.25 ^a	0.28ª	0.29 ^a	0.03
Bile	0.12 ^{ab}	0.09 ^b	0.13 ^{ab}	0.17 ^a	0.14 ^{ab}	0.14 ^{ab}	0.02
Spleen	0.14ª	0.14 ^a	0.14 ^a	0.09 ^a	0.10 ^a	0.13 ^a	0.02

Table 4. Effect of FOSPM on Internal Organ Characteristics of Broiler Chickens

^{a,b,c,} Means with different superscripts in the same row are significantly different (p<0.05), FSOPM, fermented sweet orange peel meal , SEM, standard error of mean

Parameters (%)	Treatments						
	0%	2%	4%	6%	8%	10%	
	FSOPM	FSOPM	FSOPM	FSOPM	FSOPM	FSOPM	
Live weight (g)	2.00 ^a	1.83 ^{abc}	1.77 ^{abc}	1.95 ^{ab}	1.73 ^{bc}	1.59°	0.08
GIT weight (g)	57.05 ^b	67.59 ^a	68.19 ^a	72.96 ^a	62.08 ^b	70.74 ^a	1.65
Small intestine (% GIT weight)	81.02ª	83.50 ^a	78.68 ^a	80.76 ^a	81.65ª	82.24ª	1.91
Large intestine (% GIT weight)	9.19 ^{ab}	6.24 ^b	13.63 ^a	10.39 ^{ab}	8.16 ^b	7.80 ^b	1.58
Caecum (% GIT weight)	7.04 ^a	9.96 ^a	7.72 ^a	8.98 ^a	10.37ª	10.07ª	1.57
GIT length (cm)	227.66ª	258.33ª	255.93ª	261.83ª	243.93ª	253.66ª	1.80
Small intestine (% GIT length)	80.04 ^b	83.30 ^{ab}	83.03 ^{ab}	85.48ª	82.29 ^{ab}	83.10 ^{ab}	1.00
Large intestine (% GIT length)	5.36 ^a	4.37ª	4.76 ^a	4.86 ^a	5.27ª	5.05ª	0.31
Caecum (% GIT length)	12.89 ^a	14.49 ^a	13.74ª	15.20ª	15.37ª	15.62ª	1.26

Table 5. Effect of FSOPM on GIT Morphometry of Broiler Chickens

^{a,b,c,} Means with different superscripts in the same row are significantly different (p<0.05), FSOPM, fermented sweet orange peel meal, SEM, standard error of mean

4. CONCLUSION AND RECOMMENDA-TION

This study examined the nutrient value of FSOPM as possible replacement for maize in the diets of broiler chickens and from the results obtained, the following conclusions were drawn;

- FSOPM was found to have appreciable amount of DM, CP, CF, EE, ash, NFE and ME which can relatively be compared to maize, a conventional energy feed stuff. The anti- nutritional factors were observed to be acceptable without causing any harm to broiler chickens.
- ii. The use of FSOPM at 10% as replacement of maize in broiler chicken diets supported carcass characteristics, increased GIT weight, increased large intestine weight, increased small intestine length and recessed abdominal fat pad.

It is therefore recommended that further investigatory studies be carried out at higher inclusion levels (above 10%) of FSOPM in broiler chicken diets and carcass evaluated.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- 1. FAO. FAOSTAT. Food Production; 2020. Available:http://www.fao.org/faostat/en/#da ta/QC(accessed on 24 June 2020).
- 2. Sunmola TA, Tuleun CD and Oluremi OIA. Use of sweet orange peel meal (SOPM) in broiler chicken diet with and without polyzyme. Malaysian Journal of Animal Science. 2019;22(2):35-44.
- 3. Alshelmani MI, Loh TC, Foo HL, Sazili AQ, Lau WH. Effect of feeding different levels of palm kernel cake fermented by *Paenibacillus polymyxa* ATCC 842 on nutrient digestibility, intestinal morphology, and gut microflora in broiler chickens. Animal Feed Science and Technology. 2016;216:216-224.
- Alshelmani MI, Kaka U, Abdalla EA, Humam AM, Zamani HU. Effect of feeding fermented and non-fermented palm kernel cake on the performance of broiler chickens: a review. World's Poultry Science Journal. 2021;77(2):377-388.
- Tuleun CD, Carew SN, and Ajiji TA. Feeding Value of Velvet Beans (*Mucuna utilis*) for Laying Hens. Livestock Research for Rural Development, 2008;20: 81 – 83.

- Kairalla MA, Alshelmani MI, Aburas AA. Effect of diet supplemented with graded levels of garlic (*Allium sativum* L.) powder on growth performance, carcass characteristics, blood hematology, and biochemistry of broilers. Open Veterinary Journal. 2022;12(5):595-601.
- Kairalla MA, Aburas AA, Alshelmani MI. Effect of diet supplemented with graded levels of ginger (*Zingiber officinale*) powder on growth performance, hematological parameters, and serum lipids of broiler chickens. Archives of Razi Institute. 2022;77(6):2089-2095.
- Kairalla MA, Alshelmani MI, Imdakim MM. 8. Effect of diet supplemented with different levels of moringa powder on performance, arowth carcass characteristics, meat quality, hematological parameters. serum lipids. and economic efficiency of broiler chickens. Archives of Razi Institute. 2023:78(5):1647.
- Oluremi OIA, Shiddi J, Mahlehla M and Adenkola A. Utilization of biodegraded sweet orange (*Citrus sinensis*) Fruit Peel as a replacement for maize in Growing Rabbit Diet. East African Scholary Journal of Agriculture and Life Sciences. 2020;3(10):311 – 317.
- Akpe ME, Oluremi, OIA and Tuleun CD. Haematological and Serum Biochemical Indices of Broiler Chickens fed Diets containing Graded Levels of Biodegraded Sweet Orange (*Citrus sinensis*) Peel. Journal of Agriculture and Veterinary Science. 2019;12(8):54 – 59.
- Oluremi OlA, Shiddi J, Mahlehla M, Adenkola A. Utilization of biodegraded sweet orange (*Citrus sinensis*) fruit peel as a replacement for maize in growing rabbit diet. East African Scholary Journal of Agriculture and Life Sciences. 2020;3(10):311 – 317.
- 12. TAC. Makurdi weather elements records, makurdi metereological station. Nigerian Air Force, Makurdi, Nigeria: Tactical Air Command; 2009.
- AOAC. Association of Official Analytical Chemists. 17th Edition. William Press. Richard Virginia, USA; 2002.
- Kirk SR, Sawyer R, Egan H. Pearson's composition and analysis of foods. 9th Edition, Longman, Singapore; 1991.
- Harbone JB. Phytochemical Methods: A guide to modern techniques of plant analysis. 3rd Edition. Chapman and Hall

International, New York, USA. 1998;48-189.

- Ozyurt DO, Zturk BD, Apak R. Determination of total flavonoid content of *Urtica dioica* L. by a New Method. Adnan Menderes University, 4th AACD Congress, Turkey; 2004.
- 17. Aduku AO. Poultry processing and marketing in Nigeria. Poultry Production in Nigeria, A Training Manual. 2002;146 -152.
- Minitab. Minitab statistical software reference manual. P.C. Version Release 16. New York, Media Cybernetics; 2014.
- Odunlade TA, Oluremi OIA and Abang FBP. Effect of the nutrient composition of biodegraded sweet orange (*Citrus sinensis*) Fruit peel on the Growth Performance of Starter Broiler Chicks. Animal and Veterinary Sciences, 2020;8(5):104-109.
- 20. Idoga SA. The Feeding value of biodegraded sweet orange peel in the nutrition of broiler chickens. Masters Thesis submitted to the Department of Animal Nutrition, University of Agriculture, Makurdi, Nigeria; 2021.
- 21. Ngi J. The Nutritive Potential of Sweet Orange (*Citrus sinensis*) peel meal for goat feeding. Ph.D Thesis submitted to the Department of Animal Production, University of Agriculture, Makurdi, Nigeria; 2013.
- Orayaga KT. Effect of composite mango (*Mangifera indica*) fruit reject meal on growth performance, digestibility and economics of production of rabbits. Nigerian Journal of Animal Science. 2016;1:65 – 75.
- Egberuare TAM, Alu BE, Adua MM. Carcass characteristics of broilers fed sundried sweet orange peels supplemented with quantum blue® enzyme. Journal of Applied Agriculture Research. 2020;8(1):58 – 66.
- 24. Aduku AO. Tropical feed stuff analysis table. Department of animal science, Faculty of Agricuture, Ahmadu Bello University – Samaru, Zaria, Nigeria; 2012.
- Ojabo LD, Oluremi OIA and Uza D.V. Effect of Feeding Sun- dried Sweet Orange (*Citrus sinensis*) Fruit Peel on Pullet Chick Performance. Research Opinions in Animal and Veterinary Sciences. 2014;4(9):484 -488.
- 26. Banerjee GC. A textbook on animal husbandry. Oxford and IBH publishing company pvt L.t.d. 113-B shahpurjat Asian

Games Village Side New Delhi 110049, India. 2011;1079.

- Ahaotu EO, Okonkwo JC, Maduduike FN. Performance and carcass characteristics of starter broiler birds fed processed wild cocoyam corn meal (*Colocosia esculenta*. L.) proceedings of the first Conference of Faculty of Agriculture and Veterinary Medicine Held at imo State University, Owerri, Nigeria. December, 4-6. 2017;279-287.
- Aduku AO. Tropical feedstuff analysis table. Department of Animal Science, Ahmadu Bello University, Samaru, Nigeria; 2005.
- 29. Oluremi OIA, Andrew IA, Ngi J. Evaluation of the nutritive potential of the peels of some citrus fruit varieties as feeding stuffs in livestock porduction. Pakistan Journal of Nutrition. 2007;6(6): 653 – 656.
- Oluremi OIA, Ahile AA, Jande TF. Effect of feeding graded levels of fermented sweet orange (*Citrus sinensis*) fruit peel meal on the growth and nutrient digestibility of broiler chicken. International Journal of Environment, Agriculture and Biotechnology. 2017;2:3119 – 3123.
- Orayaga KT, Oluremi OIA, Tuleun CD and Carew SN. Carcass yield and physiological status of broiler chickens fed diets containing mango (*Mangifera indica*) fruit reject meal. Nigerian Agricultural Journal, 2021;52 (3):85 – 93.
- 32. Oluyemi JA, Roberts FA. Poultry production climates. 2nd in wet Spectrum Books Edition. Limited. 2000:244.
- Aduku AO, Olukosi JO. Animal products, processing and handling in the tropics. In: Living Book Series. 3rd Edition, Abuja. G. U. Publications. 2000;32.
- Onabanjo RS, Ojewala GS, Onunkwo DN, Adedokun OO, Ewa EU, Nzotta AO and Safiyu KK. Performance of broiler chickens fed rice milling waste based diets. Nigerian Journal of Animal Production. 2021;48(2):148 – 161.
- Elfaki MOA, Abdelatti KA, Malik HEE. Effect of dietary dried rumen content on broiler performance, plasma constituents and carcass characteristic. Global Journal of Animal Science Resource, 2014;3:264-270.
- 36. Adriani L, Hernawan E, Hidayat U. Role of sweet orange (*Citrus sinensis*) waste in lowering the meat cholesterol and fat of Padjadjaran Sheep. Indonesian Journal of

Animal and Veterinary Science. 2015;20(4):250-256.

- Alefzadeh T, Bouyeh M, Hoven R, Seidavi A, Laudadio, Tufarelli V. Effect of dietary dried orange (*Citrus sinensis*) peel powder and exogenous multi-enzymes on growth and carcass traits and ileal microflora of broiler chickens. Pakistan Journal of Zoology. 2016;48(6):1891-1897.
- Nwobodo EA, Oluremi OIA, Tuleun CD, Kaankuda FG. The Effect of bio degraded sweet orange (*Citrus sinensis*) fruit peel on the growth and economic performance of starter broiler chicks. Journal of Animal Husbandry and Dairy Science. 2020;4(1):23-29.
- Guluwa LY, Oluremi OIA. and Tion MA. Replacement value of water soaked sweet orange (*Citrus sinensis*) peel meal in broiler chicken diet. American Journal of Experimental Agriculture, 2016;10(4): 1-8.
- Jimenez Moreno E, Mateos GG. Use of dietary fibre in broilers. San Juan Del Rio, Queretaro: In Memories De La Sexta Eeunion Annual Aecacem; 2013.
- 41. Svihus B. The Gizzard: Function, Influence of diet structure and effects of nutrient availability. World Poultry Science. 2011;67(2):207-223.
- Jha R, Mishra P. Dietary fibre in poultry nutrition and their effects on nutrients utilization, performance, gut health and on the environment: A review. Journal of Animal Science and Biotechnology. 2021;12:1 – 16.
- Ahamefule KU, Obua BE, Oguike SA. Blood chemistry and haematology of weaner rabbits fed sun – dried, ensiled and fermented cassava peel based diets. Pakistan Journal of Nutrition. 2006;5:248 – 253.
- 44. Oloruntola OD, Agbede JO, Ayodele SO and Oloruntola DA. Neem, pawpaw and bamboo leaf meal dietary supplementation in broiler chickens: Effect on performance and health status. Journal of Food Biochemistry. 2018;e12723. https://doi.org/10.1111/jfbc.12723
- Vikas K, Sinha AK, Makka HPS, Becher K. Dietary roles of phytate and plytase in human nutrient. A Review. Food Chemistry. 2009;120:945 – 1005.
- 46. Jorgensen H, Zhao X, Eggum B. The influence of dietary fiber source and level on development of the gastrointestinal tract, digestibility, degree of fermentation in

the hind – gut and energy metabolism in pigs. British Journal of Nutrition. 1996:75:365-378.

47. Mourao JL, Pinheiro VM, Prafes JAM, Bessa BRJ, Ferreira LMA, Fontes CMGA, Ponter PIP. Effect of dietary dehydrated pasture and citrus pulp on the performance and meat quality of broiler chickens. Poultry Science. 8200;8(7):733-743.

 Ebrahimi A, Qotbi AAA, Seidavi AR, Laudadio V. and Tufarelli V. Effect of different levels of dried sweet orange (*Citrus sinensis*) peel on broiler chickens growth performance. Archivose Tierzucht. 2013;(56):11-17.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/127515