

## **Demographics and Pattern of Ocular Diseases in Patients with Chronic Kidney Disease at the University of Port Harcourt Teaching Hospital**

**Rachael O. Aliu<sup>1</sup>, Elizabeth A. Awoyesuku<sup>2\*</sup> and Chinyere N. Pedro-Egbe<sup>2</sup>**

<sup>1</sup>*Department of Ophthalmology, Federal Medical Centre, Yenagoa, Bayelsa State, Nigeria.*

<sup>2</sup>*Department of Ophthalmology, University of Port Harcourt Teaching Hospital, Rivers State, Nigeria.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Author ROA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author EAA managed the analyses of the study. Author CNPE managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/OR/2019/v11i130118

Editor(s):

(1) Dr. Tatsuya Mimura, Department of Ophthalmology, Tokyo Women's Medical University Medical Center East, Japan.

Reviewers:

(1) Shubha Ghonsikar Jhavar, Govt. Medical College, India.

(2) Umezurike Benedict Chidozie, Government House Clinic, Nigeria.

Complete Peer review History: <https://sdiarticle4.com/review-history/51817>

**Original Research Article**

**Received 11 July 2019**  
**Accepted 25 September 2019**  
**Published 01 October 2019**

### **ABSTRACT**

**Aims:** To determine the demographics and pattern of ocular diseases in patients with Chronic Kidney Disease at the University of Port Harcourt Teaching Hospital.

**Study Design:** This was a hospital-based cross-sectional study.

**Place and Duration of Study:** This study was carried out at the Dialysis clinic of the University of Port Harcourt Teaching Hospital, Rivers State, Nigeria from February 2013 to February 2014.

**Methodology:** The Sample size was estimated to be 85 adult patients and consecutive adult patients with chronic renal failure who are having haemodialysis were enrolled into the study. The patients who met the inclusion criteria were randomly booked at the Dialysis Clinic and had no prior knowledge of the study; hence there was no bias in case selection. Ethical approval was obtained from the institutions ethical committee. Each participant had a comprehensive ocular examination including fundus photography. Subjects needing further evaluation and treatment were referred to the Eye Clinic of the Ophthalmology Department of the University of Port Harcourt Teaching Hospital, Port Harcourt.

\*Corresponding author: E-mail: [awoyeliz@yahoo.com](mailto:awoyeliz@yahoo.com);

**Results:** A total of 170 eyes of 85 persons were examined in this study (100% coverage). A total of 30 (35.4%) females and 55 (64.6%) males were examined giving a male to female ratio of 1.8:1. Approximately half of the study subjects (54.1%) had no previous eye problem prior to developing chronic renal failure while about a quarter (n= 20, 23.5%) had refractive error and difficulty in reading near prints. After diagnosis of chronic kidney disease however a total of 40 (47%) of study subjects had visual impairment while 2(2.4%) were blind and 43(50.6%) of the study population however had normal vision.

**Conclusion:** Chronic Kidney disease is an important cause of ocular morbidity in our environment with majority of those affected being males in the third to fifth decades of life. Posterior segment disorders were the commonest disorders seen.

**Keywords:** Demographics; pattern; visual impairment; chronic kidney disease.

## 1. INTRODUCTION

The World Health Organization health report shows that intrinsic diseases of the kidney and urinary tract resulted in the death of 1 million people in 2002 thereby ranking 12<sup>th</sup> on the list of major causes of death [1]. The prevalence of impaired renal function is estimated to range between 10-20% of the adult population in most countries of the world [1].

Renal disease, especially glomerular disease, is more prevalent in Africa and seems to be of a more severe form than is found in Western countries [2]. This is due to the high prevalence of infection-related nephropathies [3,4] and non-communicable diseases [5]. Similar to other developing countries of the world, no reliable statistics are available on the prevalence of kidney diseases in Nigeria[6] but hospital-based studies put the prevalence at between 3.6% to 10.4% [7,8].

In Nigeria, the 3 commonest causes of Chronic kidney disease in adults are chronic glomerulonephritis, hypertension and diabetes mellitus [9,10,11]. While common causes in children include glomerulonephritis and posterior urethral valves, [9]. Analgesics abuse, ingestion of herbs and use of skin bleaching or lightening soaps and creams containing hydroquinone and mercury are also known preventable risk factors commonly seen in Nigeria [12-14,7]. Other causes of chronic kidney disease are polycystic kidney disease, Human Immunodeficiency Virus (HIV) /Acquired Immune Deficiency Syndrome (AIDS), obstructive nephropathy, renal carcinoma, tuberculosis, sickle-cell disease, autoimmune diseases and past episode of acute renal failure [15-18]. Chronic kidney disease tends to be asymptomatic in the early stages so patients usually present late with consequent poorer prognosis [1]. End stage renal disease is the most severe form of CKD and at this stage,

patients require Renal Replacement Therapy [RRT] in the form of haemodialysis, peritoneal dialysis or renal transplant for continued survival [19,20].

The kidneys' functions include urine production, excretion of waste products of metabolism (urea, electrolytes) and production of the hormones calcitriol (active form of Vitamin D which regulates calcium metabolism) and erythropoietin (stimulates red blood cells production by the bone marrow) as well as the enzyme renin (part of renin-angiotensin system which regulates the systemic blood pressure) [21-23]. Chronic kidney disease thus leads to systemic fluid overload, electrolyte derangements, persistent uraemia and failure of the kidney to maintain its hormone secreting functions [24,25,26]. Clinical manifestations of Chronic kidney disease include heart failure, hypertension, anaemia, severe pruritus, peripheral edema, bone pain, hiccups, reduced vision, metabolic acidosis, bleeding tendencies, muscle twitching and seizures,[27]. Fluid overload can cause lid swelling which may be isolated or associated with facial puffiness [28]. At the ESRD stage, 80% of patients will have developed secondary hypertension [29].

This study aims to elucidate the demographics and pattern of ocular diseases in patients with chronic kidney disease undergoing hemodialysis in the University of Port Harcourt Teaching Hospital.

## 2. MATERIALS AND METHODS

This was a hospital-based cross-sectional study carried out at the Dialysis clinic of the University of Port Harcourt Teaching Hospital, Rivers State, Nigeria.

The Sample size was estimated to be 85 adult patients using the equation for calculation of

sample size and consecutive adult patients with chronic renal failure who are having haemodialysis were enrolled into the study.

## 2.1 Inclusion Criteria

1. Patients aged 18 years and above attending the Dialysis clinic of the University of Port Harcourt Teaching Hospital, Port Harcourt.

## 2.2 Exclusion Criteria

1. Patients below 18 years of age even if attending the Dialysis clinic.
2. Patients who did not consent to participate in the study.

The examinations were performed before patients had haemodialysis.

Measurement of blood pressure was performed by a renal unit nurse. Administration of questionnaire and obtaining informed consent was carried out. Visual acuity testing using the Snellen literate and illiterate charts and dilatation of patients' pupils with 0.5% Tropicamide was performed. Intraocular pressure measurement was done using the Perkins MK2 applanation tonometer, direct ophthalmoscopy was done using Direct ophthalmoscope (Welch-Allen, Model number 11720) while indirect ophthalmoscopy was performed using Binocular indirect ophthalmoscope (Keeler, Model number 1945-P-1001), Fundus camera (Carl Zeiss, mydriatic. FF 450 Plus model) was used for fundus photography and at the end data was entered into spread sheets for analysis [30]. They were later fed into the database of the public domain statistical software package for epidemiology, EPI info (version 7.14) designed by the Centre for disease control and Prevention in Atlanta, Georgia (USA) and analysed with the aid of a statistician. Frequency was presented in

percentages. Mean and standard deviations were calculated for descriptive and comparative purposes. Statistical significance was tested using the chi-square test. P-value < 0.05 was taken as statistically significant.

## 3. RESULTS AND DISCUSSION

170 eyes of 85 patients were included in the study. The age and sex distribution is as follows.

The age range of patients in this study is 19 to 83 years with a mean of  $43.5 \pm 15.56$  years. This is similar to findings by Alasia et al. [11] ( $46.2 \pm 17.6$  years) in a study carried out in this centre and other studies carried out in Nigeria and other developing countries; Alebiosu et al. [17] ( $39.6 \pm 14.8$  years), Bamgboye [31] (38.6 years). Arogundade [32] ( $39.9 \pm 1.67$  years), Ulasi [33] ( $42.5 \pm 15.43$  years), Quattara et al. [34] ( $44 \pm 10$  years). This however, differs from findings in developed countries. Karras et al. [35] in a study conducted in France, showed a mean age of  $59.8 \pm 14.5$  years while a United Kingdom review of general practice computerized data revealed a mean age of  $57 \pm 18.9$  years for all stages of CKD [36]. The mean age noted in developed countries is higher which implies an older age of onset of renal failure with incidence of ESRD being higher in elderly people than in the general population [37]. Renal function deteriorates with aging [38]. The extent of age-related glomerular filtration rate (GFR) decline, however, differs between ethnic groups (blacks are more affected than whites) and sexes (males more affected than females) [38]. A total of 79 (92.9%) subjects had formal education. Of this number, 43.5% (n=37) had tertiary education and most of the educated were males (54.1%) (Fig.1).

As shown in Fig. 2, 58 (68.2%) subjects were employed. Out of those employed, 28 (32.9%) were self-employed. 16 subjects (19%) were unemployed.

**Table 1. Age and sex distribution of study subjects**

Age range (years)	Sex		Total
	Female	Male	
<20	0 (0%)	2 (2.4%)	2 (2.4%)
20-29	4 (4.7%)	9 (10.6%)	13 (15.3%)
30-39	8 (9.4%)	16 (18.8%)	24 (28.2%)
40-49	6 (7.1%)	13 (15.3%)	19 (22.4%)
50-59	6 (7.1%)	7 (8.2%)	13 (15.3%)
60-69	4 (4.7%)	4 (4.7%)	8 (9.4%)
70-79	1 (1.2%)	2 (2.3%)	3 (3.5%)
80-89	1 (1.2%)	2 (2.3%)	3 (3.5%)
Total	30 (35.4%)	55 (54.6%)	100.0%




Fig. 1. Educational level of study subjects




Fig. 2. Occupation of study population

Table 2 below shows the causes of chronic kidney disease in the sample population. The commonest cause of renal failure was chronic glomerulonephritis (n=38; 44.7%) followed by

hypertension (n=23; 27.1%) then diabetes mellitus (n=11; 12.9%). The least common causes were sickle cell disease and renal carcinoma each contributing 1.2%. Major causes

of Chronic Kidney Disease/End Stage Renal Disease tend to occur at a younger age in blacks [33]. The male to female ratio of 1.8:1 found in this study is similar to worldwide data [39]. In Nigeria, Alasia et al. [11] in Port Harcourt got a male to female ratio of 1.9; 1; Alebiosu et al. [17] in Sagamu; 1.42; 1 and Ulasi et al. [40] in Enugu; 1.9; 1. The reasons for this male preponderance are unknown but in Sub-Saharan Africa, families value male more than female members and may therefore spending more money on them for medical treatment [31]. However, it could be due to faster rate of deterioration of kidney function in males with some forms of glomerulonephritis and polycystic kidney disease [38].

**Table 2. Causes of Chronic kidney disease in 85 subjects**

Cause of chronic renal failure	Frequency	Percentage
Chronic glomerulonephritis	38	44.7%
Hypertension	23	27.1%
Diabetes Mellitus	11	12.9%
HIV Associated	6	7.1%
Nephropathy		
Obstructive nephropathy	3	3.5%
Adult polycystic kidney disease	2	2.3%
Sickle cell disease	1	1.2%
Renal carcinoma	1	1.2%
<b>Total</b>	<b>85</b>	<b>100.0%</b>

The 3 commonest causes of Chronic Kidney Disease in this study were chronic glomerulonephritis (44.7%), hypertension (27.1%) and diabetes mellitus (12.9%). This is in keeping with studies from other hospitals in Nigeria and some developing countries. [28,41,42,43]. A rise in the percentage contribution of diabetes has however been noted in various centers in Africa, which is consistent with the increase in urbanization and improvement in the living standards in these countries [44]. In Western countries however, diabetes and hypertension alone are the leading causes [2,6]. Glomerular disease is more prevalent in Africa and this is due to the high prevalence of infection-related nephropathies [3,4]. These infections include infected scabies, plasmodium malariae, schistosomiasis, mycobacterium leprae, filarial worms, toxoplasmosis and streptococcal organisms [33]. The use of skin lightening creams and herbal remedies is also prevalent in our environment [7].

**Table 3. Pattern of fundal findings in study subjects**

Fundal finding	Frequency	Percentage
Hypertensive retinopathy		
Grade II	27	31.7%
Grade III	17	20.0%
Grade IV	14	16.5%
Diabetic retinopathy		
Moderate non-prolif	1	1.2%
Severe non-prolif	4	4.7%
Proliferative	5	5.9%
HIV retinopathy	6	7.1%
CRVO	1	1.2%
Tractional retinal detachment	1	1.2%
CMV retinitis	1	1.2%
Macular edema	31	36.5%
CSMO	7	8.2%
Glaucoma	4	4.7%
ARMD	3	3.5%
Optic neuropathy	2	2.4%
Diffuse retinal edema	13	15.3%
Normal fundus	3	3.5%

*Key: Non-prolif – non-proliferative; CRVO- central retinal vein occlusion; CMV –cytomegalovirus; CSMO – clinically significant macular oedema; ARMD –age-related macular degeneration*

Study subjects had more than one fundal pathology. The commonest fundus finding was hypertensive retinopathy of different grades seen in a total of 58 (68.2%) subjects, followed by macular edema seen in 31 (36.5%) patients. Only 3 (3.5%) of the study subjects had normal fundoscopic findings.

Human immunodeficiency virus retinopathy was present in half of the patients with Human Immunodeficiency Virus Associated Nephropathy (HIVAN) and Cytomegalovirus (CMV) retinitis was observed in 1 patient. This is in keeping with worldwide prevalence of HIV retinopathy of 40-60% [45,46]. Other studies did not report HIV retinopathy or CMV retinitis. This is likely because HIVAN was not reported as a cause of Chronic Kidney Disease/End Stage Renal Disease in patients from other studies [47,48,49, 50,51]. Most (70.8%) of the people living with HIV/AIDS globally live in Sub-Saharan Africa [52].

Central retinal vein occlusion was noted in 1 patient. Hypertension is an important risk factor

[53,54]. Other studies did not report retinal vascular occlusion.

Optic neuropathy was observed in 2.4% of study subjects. This is similar to findings in another study [47]. It was however not reported by other studies [48,49,50,51].

Of the 42% with visual impairment, macular edema from diabetic retinopathy is the leading cause of visual impairment in this study being accountable for half of the cases. Only a few studies assessed visual impairment in association with fundal findings [37,48]. Other studies reported maculopathy but not macular edema specifically and may have included it with figures for diabetic retinopathy. Maculopathy (Macular edema, CSMO, ARMD) accounts for a total of 22 (52.4%) cases of visual impairment. Maculopathy was reported as a much lower cause of visual impairment (9.7% of subjects) by Bajracharya et al. [47] same with Ahmed et al. [48] who reported 17%. The difference observed may be racial [55]. The higher prevalence in this study could be due to a high prevalence of hypertensive patients in the study population as hypertension has been shown to worsen diabetic retinopathy [56].

Hypertensive retinopathy stages 3 and 4 were responsible for about a third of cases of visual impairment in this study. This differs from other studies (Bajracharya et al. [47] 15%; Ahmed et al. [48] 11%). This difference may be due to a higher prevalence of hypertension in the study population. Some of the studies only assessed the patients with fundus photograph for retinopathy and did not check for visual impairment [50,57].

Diabetic retinopathy accounted for 14.3% of patients with visual impairment. This is higher than findings in other studies (Bajracharya et al. [47] 3.8%, Vrabec et al. [49] 8%, Ahmed et al. [48] 13%) The difference observed may be due to diabetic retinopathy being more severe in blacks [55].

The other causes of visual impairment were a case each of retinal detachment secondary to proliferative diabetic retinopathy and CMV retinitis in an HIV patient. The Nepal study [47] also had a case of retinal detachment while other studies did not [48,49,50,51].

Cytomegalovirus retinitis tends to occur in advanced HIV infections. Because of its strong association with both blindness and mortality in

HIV patients CMV retinitis is universally acknowledged as the clinically most important ocular complications of AIDS [46].

## CONCLUSION

Chronic Kidney Disease in our center mainly affected males and people in the 3<sup>rd</sup> to 5<sup>th</sup> decades of life. Ocular morbidity was mainly due to posterior segment (fundal) pathologies with hypertensive retinopathy topping the list.

## CONSENT

As per international standard, patient's written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

Ethical approval was obtained from the institutions ethical committee.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. WHO. The World Health Report 2003: shaping the future. WHO; 2003.
2. Moeller S, Gioberge S, Brown G. ESRD patients in 2001. Global overview of patients, treatment modalities and development trends. *Nephrol Dial Transplant*. 2002;17:2071–2076.
3. Sumaili EK, Cohen EP, Zinga CV, Krzesinski JM, Pakasa NM, Nseka NM. High prevalence of undiagnosed chronic kidney disease among at-risk population in Kinshasa, the Democratic Republic of Congo. *Nephrol Dial Transpl*. 2009;24(1): 117-122.
4. Akinsola W, Odesanmi WO, Ogunniyi JO, Ladipo GOA. Diseases causing chronic renal failure in Nigerians—a prospective study of 100 cases. *Afr J Med Med Sci*. 1989;18(2):131–137.
5. National expert committee on communicable diseases (NCD), Non-communicable diseases. Final report of a National Survey, Federal Ministry of Health and Social Services, Lagos. 1997;1-12.
6. Bommer J. Prevalence and socio-economic aspects of chronic kidney

- disease. *Nephrol Dial Transplant*. 2002; 17(11):8–12.
7. Campbell RC, Ruggeneti P, Remuzzi G: Halting the progression of chronic nephropathy. *J Am Soc Nephrol*. 2002; 13(3):190–195.
8. Lysaght MJ. Maintenance dialysis population dynamics: Current trends and long-term implications. *J Am Soc Nephrol*. 2002;13(1):37–40.
9. Phuong-Thu T Pham, Phuong-Chi T Pham, Alan H Wilkinson, Susie Q Lew. Renal abnormalities in sickle cell disease. *Kidney International*. 2000;57:1–8.
10. Chadban S, Briganti E, Kerr P. Prevalence of kidney damage in Australian adults. The Aus Diab Kidney Study. *J Am Soc Nephrol*. 2003;14(7):131–138.
11. Alasia DD, Emem-Chioma P, Wokoma FS. A single-center 7-Year experience with end-stage renal disease care in Nigeria—A Surrogate for the poor state of ESRD care in Nigeria and Other Sub-Saharan African countries: Advocacy for a global fund for ESRD care program in Sub-Saharan African Countries. *Int J Nephrol*. 2012; 10(6):1-7.
12. Ouattara B, Kra O, Yao H, Kadjo K, Niamkey EK. Characteristics of chronic renal failure in black adult patients hospitalized in the Internal Medicine department of Treichville University Hospital. *Nephrol Ther*. 2011;7(7):531-534.
13. Diouf B, Ka EF, Niang A, Diouf ML, Mbengue M, Diop TM. Etiologies of chronic renal insufficiency in an adult internal medicine service in Dakar. *Dakar Med*. 2000;45(1):62-65.
14. Ibrahim HU, Elechi HA, Rabasa A, Ashr GM, Farouk AG, et al. Prevalence and pattern of human immunodeficiency virus-associated nephropathy among human immunodeficiency virus-positive children at the University of Maiduguri Teaching Hospital, Nigeria. *Saudi J Kidney Dis. Transpl*. 2019;30(4):843-852.
15. Adetuyibi A, Akisanya JB, Onadeko BO. Analysis of the causes of death on the medical wards of the University College Hospital, Ibadan over a 14 year period (1960–1973). *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 1976;70(5-6):466–473.
16. Wang J, Staessen JA, Fagard RH. The Systolic Hypertension in China (Syst-China) Trial collaborative group: Prognostic significance of serum creatinine and uric acid in older Chinese patients with isolated systolic hypertension. *Hypertension*. 2001;37:1069–1074.
17. Alebiosu CO, Ayodele O, Adigun A, Olutoyin AI. Chronic renal failure at the Olabisi Onabanjo University Teaching Hospital, Sagamu, Nigeria. *Afr Health Sci*. 2006;6(3):132-138.
18. Afolabi MO, Abioye-Kuteyi EA, Arogundade FA, Bello IS. Prevalence of chronic kidney disease in a Nigerian family practice population. *SA Fam Pract*. 2009; 51(2):132-137.
19. Chee Wai Wong, Tien Yin Wong, Ching-Yu Cheng, Cahrumathi Sabanayagam. Kidney and eye diseases: Common risk factors, etiological mechanisms and pathways. *Kidney International*. 2013;85:1290-1302.
20. Hsiao CH, Chao A, Chu SY. Association of severity of conjunctival and corneal calcification with all-cause 1-year mortality in maintenance haemodialysis patients. *Nephrol Dial Transplant*. 2011;26:1016.
21. Umezudike T, Mabayoje M, Okany C, Abdulkareem F, Adeyomoye A, Okubadejo N, et al. Prevalence of chronic kidney disease in HIV positive patients in Lagos, South-West Nigeria. *Nephrol Reviews*. 2012;4(1):22-23.
22. Shirelle Assaram, Nombulelop. Magula, Suman Mewa Kinoo Tivani P. Mashamba-Thompson. Renal manifestationsof HIV during the antiretroviralera inSouth Africa: A systematic scoping review. *Syst Rev*. 2017;6:200
23. Fabian J, Naicker S, Venter WD. Urinary screening abnormalities in antiretroviral-naïve HIV-infected outpatients and implications for management-a single centre study in South Africa. *Ethn Dis*. 2009;19(1):80-85.
24. Gao B, Zhu L, Pan Y, Yang S, Zhang L, Wang H. Ocular fundus pathology and Chronic kidney disease in a Chinese population. *BMC Nephrol*. 2011;17(12):62.
25. Klein R, Klein BE, Knudtson MD, Wong TY, Cotch MF Liu K, et al. Prevalence of age-related macular degeneration in 4 racial/ethnic groups in the multiethnic study of atherosclerosis. *Ophthalmology*. 2006;113(3):373-380.
26. Liew G, Mitchel P, Wong TY, Iyengar SK, Wang JJ. Chronic kidney disease increases the risk of age-related macular

- degeneration. *J Am Soc Nephrol*. 2008; 19(4):806-811
27. Skorecki K, Green J, Brenner BM. *Harrison's principles of internal medicine*. 16<sup>th</sup> edition. New York, Chicago, San Francisco. McGraw-Hill Companies, Inc. Chapter 261: Chronic Renal Failure. 2005; 1655-1660.
28. Stein JH, Hulton JJ, Kohler PO. *Internal Medicine*. 3<sup>rd</sup> ed. USA: Little Brown & Comp. 1990;809-810.
29. Arogundade FA, Sanusi AA, Hassan MO, Akinsola A. The pattern, clinical characteristics and outcome of ESRD in Ile-Ife, Nigeria: Is there a change in trend? *African Health Sciences*. 2011;11(4):594-601.
30. Update of CSN clinical practice guideline. *JASN*. 2006;17:S1-S27.
31. Bamgboye EL. Hemodialysis: Management problems in developing countries, with Nigeria as a surrogate. *Kidney Int*. 2003;63(83):93.
32. Arogundade FA, Sanusi AA, Hassan MO, Akinsola A. The pattern, clinical characteristics and outcome of ESRD in Ile-Ife, Nigeria: Is there a change in trend? *African Health Sciences*. 2011;11(4):594-601.
33. Kunitoshi I. Women live longer than men. Can this phenomenon be explained by chronic kidney disease (CKD)? Gender differences in the prevalence and incidence of CKD are discussed. *Kidney Int*. 2008;(74):415-417.
34. Alebiosu CO. Clinical diabetic nephropathy in a tropical African population. *West Afr J Med*. 2003;22(2):152-155.
35. Weatherall DJ, Ledingham JGG, Warrell DA. The epidemiology of chronic kidney disease. *Kidney Int*. 2005;67:14-18.
36. Karras A, Haymann JP, Bozec E, Metzger M, Jacquot C, Maruani et al.. Large artery stiffening and remodeling are independently associated with all-cause mortality and cardiovascular events in chronic kidney disease. *Hypertension*. 2012;60:1451-1457.
37. De Lusignea S, Chanb T, Stevensc P, O'Donoghued D, Haguea N, Dzregaha B, et al. Identifying patients with chronic kidney disease from general practice computer records. *Family Practice*. 2005; 22:234-241.
38. United States Renal Data System. Annual data report: Incidence and prevalence of ESRD (2003). *Am J Kidney Dis*. 2003; 42(5):37-173.
39. Kanski JJ, Bowling B. *Clinical Ophthalmology: A systematic approach*. 5<sup>th</sup> Edition. Elsevier Saunders. 2004;194.
40. Chijioke A, Adeniyi AB. End Stage renal disease; racial differences. *OJM*. 2003; 15(1):24-31.
41. Akinsola W, Odesanmi WO, Ogunniyi JO, Ladipo GOA. Diseases causing chronic renal failure in Nigerians—a prospective study of 100 cases. *Afr J Med Med Sci*. 1989;18(2):131-137.
42. National Expert Committee on Communicable Diseases (NCD), Non-communicable diseases. Final report of a National Survey, Federal Ministry of Health and Social Services, Lagos. 1997;1-12.
43. Naicker S. End-stage renal disease in sub-Saharan and South Africa. *Kidney Int Suppl*. 2003;83:119-122.
44. Plange-Rhule J, Phillips R, Achaempong JW, Sagger-Malik AK, Cappuccio FP, Eastwood JB. Hypertension and renal failure in Kumasi Ghana. *J Hum Hypertens*. 1999;13(1):37-40.
45. Kestelyn PG, Cunningham ET Jr. HIV/AIDS and blindness. *Bull World Health Organ*. 2001;79:208-213.
46. Vrabec TR. Posterior segment manifestations of HIV/AIDS. *Surv Ophthalmol*. 2004;49:131-157.
47. Bajracharya L, Shah DN, Raut KB, Koirala S. Ocular evaluation in patients with chronic renal failure—a hospital based study. *Nepal Med Coll J*. 2008;10 (4):209-214.
48. Wong TY, Coresh J, Klein R, Muntner P, Couper DJ, Sharrett AR. Retinal microvascular abnormalities and renal dysfunction: The atherosclerosis risk in communities study. *J Am Soc Nephrol*. 2004;15:2469-2476.
49. Vrabec R, Vatauvuk Z, Pavlovic D. Ocular findings in patients with chronic renal failure undergoing hemodialysis. *Coll. Antropol*. 2005;29(1):95-98.
50. Deva R, Alias MA, Colville D, Tow FK, Ooi QL, Chew S, et al. Vision-threatening retinal abnormalities in chronic kidney disease stages 3 to 5. *CJASN*. 2011;6(8); 1866-1871.
51. Ahmed E. Khatatbeh MD, Mohammad N. Maryan MD, Amer T. Al Bderat MD, et al. Ophthalmic findings among Jordanian



- patients with Chronic Renal Failure on Haemodialysis at Prince Ali Military Hospital in the South of Jordan. JRMS September. 2013;20(3):62-67.
52. Jawa A, Koomi J, Fonseca V. Diabetic nephropathy and retinopathy. Medical clinics of North America. 2004;88(4):1001-1036.
  53. Kian-Ersi F, Taheri S, Akhlaghi MR. Ocular Disorders in Renal Transplant Patients. Saudi J Kid Dis Transpl. 2008;19(5):751-755.
  54. Fiebai B, Ejimadu CS, Komolafe RD. Incidence and risk factors for retinal vein occlusion at the University of Port Harcourt Teaching Hospital, Port Harcourt, Nigeria. Niger J Clin Pract. 2014;17(4):462-466.
  55. Kang ES, Tevlin MT, Wang YB. Hemodialysis hypotension: Interaction of inhibitors, iNOS, and the interdialytic period. Am J Med Sci. 1999;317:9-21.
  56. Grunwald JE, Alexander J, Maguire M, Whittock R, Parker C, McWilliams K, et al. Prevalence of ocular fundus pathology in patients with chronic kidney disease. Clin Am Soc Nephrol. 2010;10:2215-2223.
  57. Ladipo GO. Hypertensive retinopathy in Nigerians: A prospective clinical study of 350 cases. Trop Geo Med. 1981;33(4): 311-316.

© 2019 Aliu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://sdiarticle4.com/review-history/51817>