



## **Production of Kunapajala and Sanjibani, Their Nutritional Contributions, Microbial and Pesticidal Effect**

**Bishal Chakraborty<sup>1</sup>, Indrajit Sarkar<sup>1\*</sup>, Swathi Kulukunde<sup>2</sup>, Soumen Maitra<sup>1</sup>,  
Arpita Mandal Khan<sup>1</sup>, Sekhar Bandyopadhyay<sup>3</sup> and Abhas Kumar Sinha<sup>4</sup>**

<sup>1</sup>Department of Floriculture, Medicinal and Aromatic Plants, Faculty of Horticulture, Uttar Banga Krishi Viswavidyalaya, Pundibari, Coochbehar-736165, West Bengal, India.

<sup>2</sup>Agricultural Polytechnic, Jagtial, Telangana, India.

<sup>3</sup>Department of Plant Pathology, Faculty of Agriculture, Uttar Banga Krishi Viswavidyalaya, Pundibari, Coochbehar-736165, West Bengal, India.

<sup>4</sup>Department of Soil Science and Agricultural Chemistry, Faculty of Agriculture, Uttar Banga Krishi Viswavidyalaya, Pundibari, Coochbehar-736165, West Bengal, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/CJAST/2019/v37i230278

#### **Editor(s):**

(1) Dr. Sangita Sahni, Assistant Professor, Department of Plant Pathology, Tirhut College of Agriculture, Dr. Rajendra Prasad Central Agricultural University, India.

(2) Dr. Tushar Ranjan, Assistant Professor, Department of Molecular Biology and Genetic Engineering, Bihar Agricultural University, Sabour, India.

#### **Reviewers:**

(1) Amr Abdelkhalek Ahmed Mohamed, Egypt.

(2) E. Siva Rami Reddy, Tanta University, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49861>

**Original Research Article**

**Received 06 June 2019**

**Accepted 19 August 2019**

**Published 23 August 2019**

### **ABSTRACT**

A study on nutritional and microbial analysis of Kunapajala with different storage time interval was conducted in the Department of Soil Science & Agricultural Chemistry and the Department of Plant Pathology, UBKV, Coochbehar-736165, West Bengal during March, 2019. The motive of this work was to estimate the physicochemical properties, macro and micro nutrient content and various microbial load of Kunapajala with different storage time interval. Kunapajala had the highest P, K, Ca, Mg, Fe, Zn, Cu & Mn 40 days after preparation and it had highest N and S 20 days after preparation. It had the highest beneficial microbial load of Fungi, *Actinomycetes*, *Pseudomonas*,

\*Corresponding author: E-mail: [indrajitsarkar\\_kpg@yahoo.co.in](mailto:indrajitsarkar_kpg@yahoo.co.in);

Phosphorus Solubilising Bacteria (PSB), *Azotobacter*, *Azospirillum*, *Rhizobium* and *Trichoderma* 40 days after preparation. So, continuous foliar and soil application of Kunapajala from 20 days after preparation to 40 days after preparation was beneficial to get maximum utilization. Moreover, Kunapajala can be used as an alternative against chemical fertilizers and pesticides to develop organic farming.

**Keywords:** *Kunapajala; sanjibani; liquid organic manure; organic farming; organic pesticide; organic fertilizer.*

## 1. INTRODUCTION

India faced several famines in its history and these famines claimed millions of lives. In the famine of 1943, India lost around four million lives in eastern India alone [1]. To solve that situation and to become self-sufficient in food production, the government of India launched several scientific ventures. Ultimately in late 1960's, India became self-sufficient in food through green revolution. The success of green revolution mainly relied on the heavy use of chemical fertilizers, pesticides, high yielding varieties and modern mechanical agricultural instruments [2]. In contrast, modernization of agriculture and dependency on chemical fertilizers and pesticides gradually deteriorates the soil fertility and adversely affects the ecological balance, natural biodiversity and environment [3]. Adaptation of organic agriculture is the only way to solve this problem [4]. The procedure of preparation of Kunapajala was mentioned in Vrikshayurveda written by Surpala. According to verse 101, 102, 103 and 104 of Vrikshayurveda, it could be prepared by mixing excreta, marrow of the bones, flesh, brain and blood of the boar with water. After that the mixture should be boiled and stored in an iron pot after adding sufficient quantity of husk, sesame oil cake, honey, black gram and ghee. As per availability, the blood, flesh and marrow of fish, goat or other animals could be used for the preparation. The items should be taken at random, no specific proportion is mentioned. Verse 106 of Vrikshayurveda explained that Kunapajala was highly effective for the crop plants. A significant increase in production was observed due to spraying of Kunapajala in several crop plants including mango (*Mangifera indica*), Soapnut (*Sapindus emarginatus*), Coconut (*Cocos nucifera*), kiwi fruit (*Actinidia deliciosa*) and brinjal (*Solanum melongana*). Spraying of Kunapajala on tea bushes controlled the attack of tea mosquito bug (*Helopeltis theivora*) and loopers (*Biston suppressaria*). Narayanan (2006) reported that after spraying Kunapajala (made of rat flesh, Mushika kunapa)

rats were totally disappeared from tea garden. So it can also be used as an alternative against chemical pesticides and rodenticides [5,6,7,8,9] (Narayanan, 2006) (Bhat and Vasanthi 2008). Hence, my motive of this research is to observe the physical, nutritional and microbial properties of the Kunapajala with different storage time intervals. Due to several microbial interactions, the nutritional status of Kunapajala is continuously changing. By studying the nutritional content and microbial population, we can understand the potentiality of Kunapajala in different time intervals. According to that proper spraying schedule of Kunapajala should be recommended to the farmers for maximising crop yield.

## 2. MATERIALS AND METHODOLOGY

### 2.1 Preparation of Kunapajala

**Ingredient:** Bombay Duck fish [*Harpodon nehereus*, cheap, devoid of scales and easily decomposable) (2.5 kg), Powdered sesame oil cake (1 kg), Rice husk (1 kg), Molasses (1 Kg), Jersey cow urine (7.5 litres).

**Procedure:** All these ingredients were mixed in an earthen pot, closed the container and allowed them to ferment. Stirring twice in a day should be done in both directions. After 40 days the solution should be filtered and collected [10].

### 2.2 Preparation of Sanjibani

**Ingredients:** Jersey cow dung (1 kg), Jersey cow urine (1 litre) and Water (2 litres).

**Procedure:** Sanjibani solution was prepared by mixing all the above items. The preparation was kept in a separate earthen container. The mouth of the container was covered with a muslin cloth and kept in the shade. The preparation was then left for incubation for 9 days. During this incubation period this was stirred twice a day (morning and evening), 10 times of clock wise and anti-clockwise direction (Ali et al. 2011).



**Fig. 1. Kunapajala- Fermentation State and liquid extract after filtering**

### 2.3 Nutritional and Microbial Analysis of Kunapajala and Sanjibani

The physical, nutritional and biological parameters of Kunapajala were analysed on the day of preparation (0 days), 20 days after

preparation and 40 days after preparation using scientifically approved standard procedures. The standard procedures performed for the estimations of these parameters are described in Table 1 and Table 2.

**Table 1. Physical and chemical properties of Kunapajala**

Sl. No.	Parameters	Methods	Reference
1	Colour	Visual evaluation	
2	Odour	Sensory evaluation	
3	Mould Growth	Visual evaluation	
4	Maggot Population	Visual evaluation	
5	pH	pH meter method	[11]
6	EC	Conductivity meter method	[11]
7	Organic Carbon (OC)	Walkley and Black wet digestion	[12]
8	Total Nitrogen	Microkjeldhal method	[11]
9	Total Phosphorus	Nitric-Perchloric(9:4) digestion and colorimetry using vanado-molybdo phosphoric yellow colour method	[11]
10	Total Potassium	Nitric-perchloric(9:4) digestion and flame photometry	[11]
11	Total Calcium	Nitric-perchloric(9:4) digestion and AAS	[11]
12	Total Magnesium	Nitric-perchloric(9:4) digestion and AAS	[11]
13	Total Sulphur	Nitric-perchloric(9:4) digestion and Turbidimetry	[13]
14	Total Micronutrients Fe, Mn, Zn ,Cu	Nitric-perchloric(9:4) digestion and AAS	[11]

**Table 2. Biological properties of Kunapajala**

Sl. No.	Parameters	Methods	Reference
1	Bacteria	Nutrient Agar medium	[14]
2	Fungi	Martin's rose Bengal Agar	[15]
3	<i>Actinomycetes</i>	Ken knight's Agar medium	[16]
4	PSB	Pikovskaya's medium	[17]
5	<i>Azospirillum</i>	Nitrogen free Bromothymol blue medium	[18]
6	<i>Azotobacter</i>	Jensen's medium	[19]
7	<i>Trichoderma</i>	Trichoderma specific Medium	[20]
8	<i>Pseudomonas</i>	King's B Agar medium	[21]
9	<i>Rhizobium</i>	Yeast extract Mannitol Agar with Congo red	[22]

### 3. RESULTS AND DISCUSSION

The colour of freshly prepared Kunapajala was brownish orange then it became darker from the 20<sup>th</sup> day onwards. Through anaerobic respiration, several gases were produced and that cause natural liquids and liquefying tissues. They also caused build-up of pressure combined with the loss of integrity of the skin and ultimately the tissue was ruptured. Ruptures in the skin allowed oxygen to re-enter the tissue and provide more surface area for the development of fly larvae and the activity of aerobic microorganisms. For these activities dark brownish orange colour was developed [23,24].

Fresh preparation of Kunapajala possessed a foul alcoholic smell. Extreme foul odour was observed from 20 to 40 days onwards. Foul alcoholic odour was developed due to putrefaction. Anaerobic metabolism took place, leading to the accumulation of gases, such as hydrogen sulphide, carbon dioxide, methane, cadaverine, putrescine and nitrogen. The purging of gases and fluids resulted the strong distinctive odours [24,25].

Initially there was no mould growth in Kunapajala whereas it was first observed 5 days after preparation. Mould growth was observed on the liquid surface and also on the sides of the storage vessel from the 15<sup>th</sup> day onwards, the decrease in mould growth was observed in the 20<sup>th</sup> day and was completely absent in the 25<sup>th</sup>. Fungi consumed energy or food from the

decaying tissue and enhanced the decomposition process. Fungi were abundant in the environment. Through air or from any other source they might be appeared in the Kunapajala vessel, but when tissues became totally liquefied or almost decomposed their population started declining. It was due to unavailability of food from that decaying tissue [26,27,28].

During decomposition, at initial stages Kunapajala attracted flies and these flies which laid eggs on it. From those eggs maggots were developed. Young maggots spread throughout the container and took food from the decaying tissue. Due to the activity of maggots the tissue started decomposing faster and the bacterial activity also enhanced. This was the reason behind the high development of maggots in Kunapajala after 5 days of its preparation. After 25 days of its preparation due to the loss of readily available cadaveric material, maggot population drastically reduced [29,30,31,24,23].

On the day of preparation Kunapajala showed pH (6.74) and after 20 days it became highly acidic in nature (3.47). Then after 40 days it became alkaline in nature (8.81). Animal tissue decomposition initially created an alkaline environment and due to microbial activity it became acidic after 20 days. When decomposition was totally completed, it became alkaline again [32,33,34,35,36]. Similar results were also found by Anandan et al. [37,38,39] in Kunapajala.

**Table 3. Physical and physico-chemical parameters of Kunapajala and Sanjibani**

KUNAPAJALA			
Parameters	On the day of preparation (0 days)	20 days after preparation	40 days after preparation
Colour	Light brownish orange	Brownish orange	Dark brownish orange
Odour	Mild alcoholic smell	Foul alcoholic smell	Extreme foul alcoholic smell
Mould growth	No mould growth	Heavy mould growth	No mould growth
Maggot Population	No maggot found	Heavy maggot growth	No maggot found
pH	6.74	3.47	8.81
EC (ds/m)	2.55	9.72	8.57

SANJIBANI			
Parameters	On the day of preparation (0 days)	20 days after preparation	40 days after preparation
Colour	Yellow	Dark yellow	Dark yellow
Odour	Foul odour	Foul odour	Extreme foul odour
Mould growth	No mould growth	Heavy mould growth	No mould growth
Maggot Population	No maggot growth	No maggot growth	No maggot growth
pH	8.73	8.84	9.05
EC (dS/m)	5.71	7.73	8.52

Kunapajala showed highest EC 20 days after preparation (9.72 ds/m) due to high acidic nature of the solution and after that it started declining (8.57 ds/m, 40 days after preparation). On the day of preparation it showed the lowest EC (2.55 ds/m) [32,33,34,35,36]. Anandan et al. [37] and Ankad et al. [39] also concluded similar trend and results.

Total OC (organic carbon) was highest 40 days after preparation (4.18%) and on the day of preparation it showed minimum value (1.72%) in Kunapajala. In decomposition physical breakdown and biochemical transformation of complex organic molecules occurred, due to that several organic carbon compounds were synthesized [40]. This was the reason for continuous increase of OC in Kunapajala. Anandan et al. [37] noticed similar trend of OC and results in his experiment.

Physical and physicochemical parameters of Kunapajala were mentioned in Table 3.

The highest N content was recorded 20 days after preparation in Kunapajala (7238 mg/dm<sup>3</sup>) while on the day of preparation it recorded the lowest value (3486 mg/dm<sup>3</sup>). For the activity of bacteria and maggots, Kunapajala started decomposing faster and due to that N content of the Kunapajala was in an increasing trend, but after 20 days 9-44% of the N was volatilized in the form of Ammonia from the solution due to alkalinity of the Kunapajala solution at that moment [41]. Ankad et al. [39] and Jani et al. [38] also concluded similar trend and results in their experiment.

On the day of preparation Kunapajala recorded the lowest value (208.661 mg/dm<sup>3</sup>) of P, while after 40 days of preparation it recorded the highest value (517.717 mg/dm<sup>3</sup>) of P. Kunapajala contained animal tissue which had high P content. According to Tian et al. [42] organic matters high in P decompose faster and release P significantly. So, Kunapajala had increasing trend of P content during decomposition. Ankad et al. [39] and Jani et al. [38] also analysed the P content of Kunapajala and found similar results.

K content was lowest on the day of preparation (890.396 mg/dm<sup>3</sup>), after that it was gradually increased and reached the highest value 40 days after preparation (1873.543 mg/dm<sup>3</sup>). Activity of

fungi and other microorganisms was the reason behind continuous release of K up to 40 days [43].

The highest Ca content was observed 40 days after preparation (614 mg/l) and on the day of preparation it was the lowest (376 mg/l). Excessive fungus and microbial activity was the reason for continuous release of Ca up to 40 days [44].

On the day of preparation Mg content was the lowest (56 mg/l) whereas after 40 days it recorded the highest value (88 mg/l). Fungal and microbial activity was the main cause behind gradual release of Mg in Kunapajala [44].

S content was the lowest on the on the day of preparation (678 mg/l), whilst 20 days it recorded the highest value (857 mg/l), then S content started declining. Due to excessive volatile release of hydrogen sulphide, after 20 days S content started declining [44].

The highest Fe content was recorded 40 days after preparation (72 mg/l) while on the day of preparation it was the lowest (55 mg/l). Due to fungal and bacterial activity gradual release of Fe was noticed in Kunapajala [45].

On the day of preparation Zn content was minimum (6.78 mg/l) while 40 days after preparation it became maximum (17.75 mg/l). Gradual increase of Zinc content was noticed in Kunapajala due to activity of fungi and bacteria [46,47,48].

Cu content was maximum 40 days after preparation (8.53 mg/l) and on the day of preparation it recorded the lowest value (4.76 mg/l). Continuously increasing trend of Cu content was observed due to activity of several fungal and bacterial species [46,47,48].

The highest Mn content was noticed 40 days after preparation (2.06 mg/l) and on the day of preparation the Mn content recorded the lowest value (0.58 mg/l). Heavy microbial interaction or activity inside Kunapajala might be the reason of this trend and result.

The macro and micro nutrient content of Kunapajala was mentioned in Table 4.

**Table 4. Macro and micro nutrient content of Kunapajala**

<b>KUNAPAJALA</b>			
<b>Parameters</b>	<b>On the day of preparation (0 days)</b>	<b>20 days after preparation</b>	<b>40 days after preparation</b>
N mg/dm <sup>3</sup>	3486	7238	4690
P mg/dm <sup>3</sup>	208.661	296.260	517.717
K mg/dm <sup>3</sup>	890.396	1589.994	1873.543
Ca (mg/l)	376	452	614
Mg (mg/l)	56	73	88
S (mg/l)	678	857	719
Fe (mg/l)	55	67	72
Zn (mg/l)	6.78	13.63	17.75
Cu (mg/l)	4.76	7.44	8.53
Mn (mg/l)	0.58	1.27	2.06

<b>SANJIBANI</b>			
<b>Parameters</b>	<b>On the day of preparation (0 days)</b>	<b>20 days after preparation</b>	<b>40 days after preparation</b>
N mg/dm <sup>3</sup>	1540.95	8358.28	7358.42
P mg/dm <sup>3</sup>	68.898	284.449	195.886
K mg/dm <sup>3</sup>	503.498	1147.445	1741.043
Ca (mg/l)	48.42	58.28	107.82
Mg (mg/l)	5.15	16.78	22.23
S (mg/l)	271.48	496.13	364.25
Fe (mg/l)	4.47	6.59	11.57
Zn (mg/l)	0.032	0.246	0.764
Cu (mg/l)	0.874	1.12	2.23
Mn (mg/l)	0.068	0.112	0.184

Fungi population was the highest 40 days after preparation ( $33 \times 10^8$  cfu/ml) and it was the lowest on the day of preparation ( $4 \times 10^4$  cfu/ml). This gradual increasing trend was noticed due to enhanced activity of early stage fungi *ascomycetes*, *deuteromycetes* and saprophytic *basidiomycetes* and late stage fungi ectomycorrhizal *basidiomycetes* in Kunapajala with time [43].

On the day of preparation Kunapajala recorded the lowest *Actinomycetes* population ( $3 \times 10^3$  cfu/ml). After that it increased continuously and reached the highest 40 days after preparation ( $5 \times 10^8$  cfu/ml). Continuous decomposition of complex mixture of polymers in dead animal tissues was the prime reason for continuous development of *Actinomycetes* population in Kunapajala [49,50,51].

The highest population of *Pseudomonas* was noticed 40 days after preparation ( $13 \times 10^{10}$  cfu/ml) in Kunapajala while on the day of preparation it recorded the lowest ( $5 \times 10^3$  cfu/ml). This type of increasing trend up to 40 days in Kunapajala was also concluded by Ali [52].

PSB population was the highest on the day of preparation ( $2 \times 10^5$  cfu/ml) then it became maximum at 40 days after preparation ( $21 \times 10^{10}$  cfu/ml) in Kunapajala. Similar trend of population growth was also observed by Ali [52] in Kunapajala.

On the day of preparation *Azotobacter* population had the lowest value ( $7 \times 10^4$  cfu/ml) in Kunapajala while after 40 days it became the highest ( $13 \times 10^{12}$  cfu/ml). Presence of *Azotobacter* in Kunapajala and this type of growth trend was justified by Ali [52].

The highest *Azospirillum* population was noticed 40 days after preparation ( $13 \times 10^{10}$  cfu/ml) and on the day of preparation the lowest value was found ( $11 \times 10^3$  cfu/ml). Ali [52] approved the existence and growth behaviour of *Azospirillum* in Kunapajala.

The lowest *Rhizobium* Population was found on the day of preparation ( $2 \times 10^3$  cfu/ml) and after 40 days, the highest *Rhizobium* population ( $4 \times 10^{11}$  cfu/ml) was noticed in Kunapajala. Ali et al. [52] also concluded similar trend of population growth of *Rhizobium* in Kunapajala.

*Trichoderma* population was highest 40 days after preparation ( $21 \times 10^8$  cfu/ml) in Kunapajala and on the day of preparation it had the lowest population ( $6 \times 10^3$  cfu/ml). *Trichoderma* had significant contribution in decomposition and biodegradation of organic

matters and due to that the population of *Trichoderma* in Kunapajala had a continuous increasing trend up to 40 days [53].


Microbial population of Kunapajala was mentioned in Table 5.

**Table 5. Microbial population of Kunapajala**

KUNAPAJALA			
Parameters	On the day of preparation (0 days)	20 days after preparation	40 days after preparation
Fungi (cfu/ml)	$4 \times 10^4$	$16 \times 10^7$	$33 \times 10^8$
<i>Actinomyces</i> (cfu/ml)	$3 \times 10^3$	$6 \times 10^4$	$5 \times 10^8$
<i>Pseudomonas</i> (cfu/ml)	$5 \times 10^3$	$8 \times 10^{10}$	$13 \times 10^{10}$
PSB(cfu/ml)	$2 \times 10^5$	$15 \times 10^{10}$	$21 \times 10^{10}$
<i>Azotobacter</i> (cfu/ml)	$7 \times 10^4$	$9 \times 10^{12}$	$13 \times 10^{12}$
<i>Azospirillum</i> (cfu/ml)	$11 \times 10^3$	$8 \times 10^8$	$13 \times 10^{10}$
<i>Rhizobium</i> (cfu/ml)	$2 \times 10^3$	$6 \times 10^6$	$4 \times 10^{11}$
<i>Trichoderma</i> (cfu/ml)	$6 \times 10^3$	$18 \times 10^8$	$21 \times 10^8$

SANJIBANI			
Parameters	On the day of preparation (0 days)	20 days after preparation	40 days after preparation
Fungi (cfu/ml)	$4 \times 10^3$	$14 \times 10^3$	$19 \times 10^3$
<i>Actinomyces</i> (cfu/ml)	0	$3 \times 10^3$	$6 \times 10^5$
<i>Pseudomonas</i> (cfu/ml)	$3 \times 10^3$	$7 \times 10^4$	$17 \times 10^4$
PSB(cfu/ml)	$3 \times 10^3$	$12 \times 10^5$	$19 \times 10^5$
<i>Azotobacter</i> (cfu/ml)	$5 \times 10^3$	$16 \times 10^5$	$13 \times 10^6$
<i>Azospirillum</i> (cfu/ml)	$2 \times 10^2$	$4 \times 10^3$	$7 \times 10^4$
<i>Rhizobium</i> (cfu/ml)	$1 \times 10^2$	$3 \times 10^3$	$6 \times 10^4$
<i>Trichoderma</i> (cfu/ml)	$3 \times 10^3$	$11 \times 10^5$	$18 \times 10^5$








Fig. 2.

#### 4. CONCLUSION

The study concludes that Kunapajala has high nutrient content and beneficial microbial population. Nutrient content of Kunapajala is highly influenced by its microbial population. Fungi will help to breakdown complex organic compounds and produce simple organic and inorganic compounds useful for plants. *Azotobacter*, *Azospirillum* and *Rhizobium* help to fix more N in crop field. PSB enhance the P solubilisation in crop field. *Actinomycetes* help to decompose complex organic molecules and antagonistic potential of *Pseudomonas*, while *Trichoderma* will help to protect the crop from soil-borne diseases. Microbial population is continuously increasing and it became the highest after 40 days. So, application of Kunapajala after 40 days is beneficial for crops, but N and S content of Kunapajala is the highest 20 days after preparation, so to exploit that spraying of Kunapajala after 20 days is also recommended. Spraying of Kunapajala on the day of preparation is not recommended because the microbial population and nutrient content is the minimum and most of the organic matter is not properly decomposed, so they will not be highly available for the crop plants. So foliar and soil application of Kunapajala from 20 days of its preparation to 40 days of its preparation is recommended for the crop and soil because we can utilise its total potential. The ingredients required to prepare it are easily available and cheap comparing with chemical fertilizers and pesticides. The crops produced using Kunapajala

will be free from any harmful chemical residues. So, it is healthy for the consumer. Moreover use of Kunapajala instead of chemical fertilizer and pesticide is highly useful to increase the crop yield, soil productivity and farmer's income.

#### ACKNOWLEDGEMENT

My special thanks to Saddam da, Salim da, Amar da and my classmate Hiralal Mandi and Somesa Ghosh for their cooperation during laboratory work.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Dyson T, Maharatna A. Excess mortality during the Bengal famine: A re-evaluation. *Indian Economic and Social History Review*. 1991;281-297.
2. FAO. Rapid growth of selected Asian economies; 2009. Available: <http://www.fao.org/docrep/009/ag087e/AG087E05.htm>
3. Paull J. Nanomaterials in food and agriculture: The big issue of small matter for organic food and farming. In: 3<sup>rd</sup> Scientific Conference of International Society of Organic Agriculture Research, Namyangju, Korea. 2011;2:96-99.



4. Manna MC, Swarup A, Wanjari RH, Ravankar HN, Mishra B, Saha MN, Singh YV, Sahi DK, Sarap PA. Long-term effect of fertilizer and manure application on soil organic carbon storage, soil quality and yield sustainability under sub-humid and semi-arid tropical India. *Fields Crop Research*. 2005;93(2-3):264-280.
5. Ayangarya VS. Herbal kunapa. *Asian Agri-History*. 2004a;8:315–317.
6. Ayangarya VS. Manujala: A liquid manure. *Asian Agri-History*. 2004b;8:319–321.
7. Ayangarya VS. INDSAFARI – An organic pesticide for tea. *Asian Agri-History*. 2005;9:317–319.
8. Ayangarya VS. Mushika kunapa. *Asian Agri-History*. 2006a;10:157–159.
9. Ayangarya VS. Kiwifruit plant treatment on the Himalayas of India: A Vrikshayurveda experience. In: *Bridging Gap between Ancient and Modern Technologies to Increase Agricultural Productivity: Proceedings of the National Conference held from 16-18 December 2005, Central Arid Zone Research Institute, Jodhpur 342 003, Rajasthan, India.* (Choudhary SL, Saxena RC, Nene YL eds.). *Asian Agri History Foundation, (AAHF), Secunderabad, India; and Rajasthan Chapter of AAHF, Udaipur, India*. 2006b;102-103.
10. Sarkar S, Kundu SS, Ghorai D. Validation of ancient liquid organics- Panchagavya and Kunapajala as plant growth promoters. *Indian Journal of Traditional Knowledge*. 2014;13(2):398-403.
11. Jackson ML. *Soil chemical analysis*. (Prentice Hall of India Pvt. Ltd, New Delhi). 1973;183-204.
12. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci*. 1934;37(1):29-38.
13. Massoumi A, Cornfield AH. A rapid method for determining sulphate in water extracts of soils. *The Analyst*. 1963;88:321–322.
14. Atlas RM, Parks LC. *Handbook of microbiological media*, CRC Press, Inc. London. 1993;529.
15. Martin JP. Use of acid, rose Bengal, and streptomycin in the plate method for estimation soil fungi. *Soil Sci*. 1950;69(3):215-232.
16. Cappuccino JG, Sheman N. *Microbiology – A laboratory manual*. 4<sup>th</sup> ed. The Benjamin/ Cummings Publishing Company. Inc., Menlo Park, California. 1996;13-182.
17. Sundararao WVB. Phosphate dissolving organisms in the soil and the rhizosphere. *Indian J. Agr. Sci*. 1963;33:272-278.
18. Dobereiner J, Marriel IE, Nery M. Ecological distribution of *Spirillum lipoferum*, Beijerinck. *Can. J. Microbiol*. 1976;22:1464-1473.
19. Jensen HL. Nitrogen fixation in leguminous plants. General characteristics of root nodule bacteria isolated from species of *Medicago* and *Trifolium* in Australia. *Proc. Linn. Soc. N.S.W.* 1942;66:98-108.
20. Saha DK, Pan S. Qualitative evaluation of some media of *Trichoderma* and *Gliocladium* spp. *J. Mycopathol. Res*. 1997;35:7-14.
21. King EO, Ward MK, Raney DE. Two simple media for the demonstration of payociamin and fluorescein. *J. Lab. Clin. Med*. 1954;44:301-307.
22. Fred EB, Baldwin IL, McCoy F. *Root nodule bacteria and leguminous plants*. University of Wisconsin Press, Madison, Wisconsin; 1932.
23. Janaway RC, Percival SL, Wilson AS. Decomposition of human remains. In Percival SL. (ed.). *Microbiology and Aging*. Springer Science + Business. 2009;13–334.
24. Carter DO, Tibbett M. Cadaver decomposition and soil: Processes. In Tibbett M, Carter DO. (eds.). *Soil Analysis in Forensic Taphonomy*, CRC Press. 2008;29–51.
25. Payne JA. A summer carrion study of the baby pig (*Sus scrofa* Linnaeus). *Ecology*. 1965;46(5):592–602.
26. Hawksworth DL, Wiltshire PEJ. Forensic mycology: The use of fungi in criminal investigations. *Forensic Sci Int*. 2011;206(1-3):1-11.
27. Schwarz P, Dannaoui E, Gehl A, Felksee-Zech H, Birngruber CG, Dettmeyer RB, Verhoff MA. Molecular identification of fungi found on decomposed human bodies

- in forensic autopsy cases. *Int J Legal Med.* 2015;129:785-791.
28. Hitosugi M, Ishii K, Yaguchi T, Chigusa Y, Kurasa A, Kido M, Nagai T, Tokudome S. Fungi can be a useful forensic tool. *Leg Med.* 2006;8:240-242.
29. Anderson GS. Minimum and maximum development rates of some forensically important Calliphoridae (Diptera). *Journal of Forensic Sciences.* 2000;45:824-832.
30. Fuller ME. The insect inhabitants of carrion: A study in animal ecology. Council for Scientific and Industrial Research, Bulletin No.-82. 1934;63.
31. Morovic-Budak A. Experiences in the process of putrefaction in corpses in buried in earth. *Medicine, Science and the Law.* 1965;5(1):40-43.
32. Carter D. Forensic taphonomy: Processes associated with cadaver decomposition in soil. Ph.D. thesis, James Cook University, Townsville, Australia; 2005.
33. Hopkins DW, Wiltshire PEJ, Turner BD. Microbial characteristics of soils from graves: An investigation at the interface of soil microbiology and forensic science. *Appl. Soil Ecol.* 2000;14:283-288.
34. Rodriguez WC, Bass WM. Decomposition of buried bodies and methods that may aid in their location. *J. Forensic Sci.* 1985;30:836-852.
35. Gill-King H. Chemical and ultrastructural aspects of decomposition. *Forensic Taphonomy: The Postmortem Fate of Human Remains.* 1997;93-108.
36. Towne EG. Prairie vegetation and soil nutrient responses to ungulate carcasses. *Oecologia.* 2000;122(2):232-239.
37. Anandan R, Priya L, Rajendran P. Dynamics of organic biofertilizers on *Oryza sativa* ADT-43. *Int. J. Curr. Microbiol. App. Sci.* 2016;5(4):902-908.
38. Jani S, Prajapati PK, Harisha CR, Patel BR. Kunapajala liquid organic manure: Preparation and its quality parameters. *World Journal of Pharmacy and Pharmaceutical Sciences.* 2017;6:1989-2000.
39. Ankad GM, Hiremath J, Patil RT, Pramod HJ. Nutrient analysis of kunapajala and panchagavya and their evaluation on germination of ashwagandha (*Withania somnifera* Dunal.) and Kalamegha (*Andrographis paniculata* Nees) seeds: A comparative study. *Journal of Ayurveda and Integrative Medicine.* 2017;xxx:1-7.
40. Juma NG. The pedosphere and its dynamics: A systems approach to soil science. Edmonton, Canada, Quality Color Press Inc. 1998;1:315.
41. Kirchmann H, Witter E. Ammonia volatilization during aerobic and anaerobic manure decomposition. *Plant and Soil.* 1989;115(1):35-41.
42. Tian G, Brussaard L, Kang BT. An index for assessing the quality of plant residues and evaluating their effects on soil and crop in the (sub-) humid tropics. *Applied Soil Ecology.* 1995;2(1):25-32.
43. Carter DO, Tibbett M. *Taphonomic mycota*: Fungi with forensic potential. *Journal of Forensic Sciences.* 2003;48(1):1-4.
44. Carter DO, Yellowlees D, Tibbett M. Cadaver decomposition in terrestrial ecosystems. *Naturwissenschaften.* 2007;94(1):12-24.
45. Dent BB, Forbes SL, Stuart BH. Review of human decomposition processes in soil. *Environmental Geology.* 2004;45(4):576-585.
46. Hodson ME, Valsami-jones E, Cotterhowells JD, Dubbin WE, Kemp AJ. Effect of bone meal (*Calcium phosphate*) amendments on metal release from contaminated soils - A leaching column study. *Environmental Pollution.* 2001;112:233-243.
47. Kearney T. Remediation of toxic metal pollution in soil using bone meal amendments. Environment Agency; 2000.
48. Deydier E, Guilet R, Sarda S, Sharrock P. Physical and chemical characterisation of crude meat and bone meal combustion residue: "Waste or raw material?" *Journal of Hazardous Materials.* 2005;121(1-3):141-148.
49. Goodfellow M, Williams ST. Ecology of actinomycetes. *Annual Review of Microbiol.* 1983;37:189-216.
50. McCarthy AJ, Williams ST. Actinomycetes as agents of biodegradation in the environment- A review. *Gene.* 1992;115:189-192.
51. Stach JE, Bull AT. Estimating and comparing the diversity of marine

- Actinobacteria. Antonie van Leeuwenhoek. 2005;87:3-9.
52. Ali MN. Sustainable agriculture with low cost technologies (SALoCT). A project funded by Rural Technology action Group – Eastern India (RuTAG-EI), IIT Kharagpur, under DST, Govt. of India; 2012.
53. Woo SL, Ruocco M, Vinale F, Nigro M, Marra R, Lombardi N. Trichoderma based products and their use in Agriculture. The open Mycology Journal. 2014;8(1):71-126.

© 2019 Chakraborty et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/49861>