



Floating and Stability Effect on Fish Feed Pellets Using Different Concentration of Baobab Leaf Meal (*Adansonia digitata*)

Eze Felix^{1*} and Eyo Victor Oscar¹

¹*Department of Fisheries and Aquaculture, Faculty of Marine Environmental Management, Nigeria Maritime University, Okerenkoko, Delta State, Nigeria.*

Authors' contributions

This work was carried out in collaboration between both authors. Author EF designed the study, wrote the protocol, performed the statistical analysis and wrote the first draft of the manuscript. Author EVO managed the analysis and literature review of the study. Both authors read and approved the manuscript.

Article Information

DOI: 10.9734/AJFAR/2018/v1i4348

Editor(s):

(1) Dr. Jorge Castro Mejia, Department of El Hombre Y Su Ambiente, Universidad Autonoma Metropolitana Xochimilco, Mexico.

(2) Dr. Matheus Ramalho de Lima, Professor, Federal University of South of Bahia, Brazil.

(3) Dr. Telat Yanik, Professor, Department of Aquaculture, Faculty of Fisheries, Ataturk University, Turkey.

Reviewers:

(1) Ali Türker, Muğla University, Turkey.

(2) Magdy Mohamed Aly Gaber, National Institute of Oceanography and Fisheries, Egypt.

(3) P. Chisoro, Wageningen University and Research, The Netherlands.

Complete Peer review History: <http://www.sciencedomain.org/review-history/26403>

Original Research Article

Received 26 June 2018

Accepted 14 September 2018

Published 27 September 2018

ABSTRACT

The objective of this study was to evaluate the floating and stability of fish feed pellets formulated with different concentration of Baobab (*Adansonia digitata*) leaf meal (BLM). Five isonitrogenous fish feed (35% CP) with varying inclusion levels of Baobab leaf meal (0% BLM, 4% BLM, 8% BLM, 12% BLM and 16% BLM designated as D₁, D₂, D₃, D₄ and D₅ respectively) was used. Ten (10) pellets of each experimental feed were placed in plastic beaker (55 x 25 x 30 cm) for 50 minutes. Feed pellet floatability and stability were recorded every five minutes. Results showed that feed pellet floatability increased significantly ($P < 0.05$) with increase in BLM concentration. Highest feed pellet floatability (41.66 ± 2.88 minutes) was recorded in D₅ (16% BLM) and least (0.00 ± 0.00 minutes) in D₁ (0% BLM) which sank down immediately. Similarly, feed pellet stability increased significantly ($P < 0.05$) with increase in BLM concentration. Highest pellet stability (42.66 ± 1.17

*Corresponding author: Email: felixeze8@gmail.com;

minutes) was recorded for D₅ (16% BLM) and least (18.54 ± 2.10 minutes) in D₁ (0% BLM). Based on these findings, it is concluded that 16% BLM inclusion level in fish feed has led to a high pellet floatability and stability. Therefore, Baobab leaf meal (BLM) which is relatively cheap, toxic free, easy to process and available specifically in Northern part of Nigeria is recommended for floating feed formulation.

Keywords: Water stability; floatability; Baobab Leaf Meal (BLM); fish feed pellets; *Adansonia digitata*.

1. INTRODUCTION

In Nigeria, fish farming is increasingly becoming very lucrative because Nigeria is one of the largest fish consumers in the world. For optimal health, fast growth and sustainable production of farmed fish, a balanced feed with a good physical characteristic such as pellet stability and floatability is required. Fish nutrition is therefore critical to sustainable aquaculture production as it represents about 60 – 80% of the total production cost [1]. According to Ilim and Cuzon, [2], aquafeed can either be pelleted or extruded with particles of high durability to withstand handling, transportation stress. High quality aquafeed should be highly stable in water to minimize disintegration and loss of nutrients upon exposure to water. Floating feed is very suitable for pelagic or surface feeders because fish quickly get access to the feed and do not expend much energy in swimming to the bottom to source for food [3]. Impaired growth has been documented on feeding fish with non-floating and unstable feed due to disintegration and sinking of feed into mud or pond bottom restricting utilization by the target fish [4]. Such disintegration may lead to bacterial build up which is capable of causing diseases to the fish. Use of stable and floating feed will help in complete utilization by the fish and minimum wastage which will help in a more profitable and sustainable aquaculture production [5,6]. Moreover, floating fish feed will enable the farmer to observe how much and how active their fish are responding to feed [7]. Baobab is a deciduous tree with a lifespan of hundreds to thousands of years [8]. Baobab spends only 4 months of the year in leaf with the fresh young leaves containing nutrients such as protein (4%), vitamin a and c [9]. The fruit pulp has a very high content of vitamin c which is almost ten (10) times that of oranges [10,11].

Baobab leaf is an excellent source of iron, calcium, potassium, manganese, molybdenum, magnesium, zinc and phosphorus. Energy value

varies from 1180-1900 kJ/100 g of which 80% is metabolized energy. The leaves are rich in pro-vitamins A and C. In terms of protein content, baobab leaves are rich in 5 out of the 8 essential amino acids [10]. In Nigeria, baobab is specifically available in the northern part of the country. Generally, baobab is comprised of eight (8) species with large, spectacular and nocturnal flowers [12]. *Adansonia digitata* is a baobab species that is indigenous to drier part of Africa while *Adansonia ghibbosa* is restricted to the North-Western Australia. The remaining six (6) species are endemic to Madagascar [13]. *A. Digitata* which grows in the arid and semi-arid region of Africa is commonly known as monkey bread which is derived from the fact that monkeys eat baobab fruit. In fish culture, one of the major factors that negatively influence growth and good health of fish is quality of feed with regards to nutrient profile and physical characteristics. Since commercial feed is highly expensive and sometimes, its availability and supply is inconsistent especially in rural areas where most fish farms are located, most farmers are presently formulating local feed using locally available ingredients [14]. The major challenges most of these fish farmers are facing is the sinking and poor stability of these locally formulated feed, which results in leaching of nutrients into the water, disintegration of feed, water pollution and growth of harmful bacteria which may predispose fish to diseases. This may result in poor growth performance of fish and reduced profitability [14]. Therefore, the objective of the study is to evaluate the effect of baobab leaf meal (*A. Digitata*) on the floatability and stability of fish feed pellets formulated with locally available raw materials.

2. METHODS

2.1 Study Area

The study was carried out in Fish Nutrition Laboratory of the department of fisheries, University of Maiduguri, Borno state, Nigeria. It is geographically located at latitude 11.80°N and

longitude 13.19°e and it is situated at elevation 325 meters above sea level.

2.2 Collection and Processing of Baobab Leaves (*A. digitata*)

Fresh baobab leaves (*A. digitata*) were collected from the Botanical garden of the University of Maiduguri, and identified by a Botanist from University of Maiduguri. The leaves were soaked in water for 24 hours in order to eliminate anti-nutritional factors. Thereafter, the leaves were sundried before grounded into powder using the hammer miller and kept in an airtight container until required.

2.3 Formulation of Experimental Diets

Five isonitrogenous fish feed (35% cp) with varying inclusion levels of Baobab Leaf Meal (0% BLM, 4% BLM, 8% BLM, 12% BLM and 16% BLM designated as d₁, d₂, d₃, d₄ and d₅ respectively) was used (Table 1). The experimental diets were formulated according to Pearson square methods.

2.4 Feed Pellet Floatability and Water Stability Test

Ten (10) pellets of each experimental feed were placed gently on the surface of water in a plastic basin of size 55 x 25 cm for 50 minutes and floatability was recorded after every 5 minutes interval. Water stability test was conducted using 10 pellets (2 mm) diameter tied in a nylon sieve material of (0.1 mm mesh). They were carefully tied with a twine to avoid breakage. Ten (10) for each treatment were fixed in a plastic basin of size 55 x 25cm and allowed to remain for time interval ranging from 10 minutes to 50 minutes with removal after every 10 minutes. At the end of every test, one of the samples for each replicate was lifted slowly with the aid of the twine and allowed to drain for 3 minutes after which the contents were put on flat boards and oven-dried at 105°C for 24 hours to obtain the whole pellet at the start of the test. The water stability (ws) was calculated using the equation below;

Water stability (mins) = weight of retained whole pellets/ initial weight of pellets x time taken.

2.5 Proximate Composition of Experimental Diets

Proximate composition of each experimental diet was analyzed according to the methods of AOAC

[15]. Protein and lipid were determined by the micro kjeldahl and soxhlet extraction of samples.

2.6 Statistical Analysis

Data obtained from the experiment were subjected to one way analysis of variance (ANOVA) with the aid of statistix version 8.0 and means separation between the treatments was done using LSD at 0.05% confidence level (p= 0.05).

3. RESULTS AND DISCUSSION

The proximate composition (Table 2) of the experimental diets formulated with varying levels of baobab leaf meal (BLM) showed that the highest crude protein (32.80%), crude fibre (19.66%), crude fat (8.10%), crude ash (2.33%) and NFE (35.95%), were obtained in d₅ (16% BLM), while the lowest crude protein (29.47%), crude fibre (13.33), crude fat (6.55%) and the highest moisture content (10.56%) were analyzed from d₁ (control). D₃ (8% BLM) had the highest dry matter of 97.30% and the lowest crude ash of (1.66%). There was no significant difference (p>0.05) between the proximate compositions of the diets with varying inclusion levels of baobab leaf meal (BLM) because the diets are isonitrogenous and the crude protein is the same.

Results obtained for the floatability of the experimental diets (Table 3) formulated with varying levels of Baobab leaf meal (BLM) showed that after 50 minutes of exposure to water, the control diet did not float at all recording a mean floating time of 0.00 ± 0.00 minutes. Furthermore, feed D₂, D₃, D₄ and D₅ showed a significant (p<0.05) improvement in their floating ability compared to the control feed (D₁). D₅ had the maximum floatation period of 41.66 ± 2.88 minutes, followed by D₄ (25.00 ± 0.00 minutes), D₃ (10.00 ± 5.00 minutes) and D₂ (8.33 ± 2.88 minutes). Results obtained for the stability of the experimental diets (Table 3) formulated with varying levels of baobab leaf meal (BLM) showed that feed formulated with BLM had a significantly higher (p<0.05) stability compared to the control feed (D₁). After 50 minutes of exposure to water, feed D₅ had the highest water stability of 42.66 ± 1.17 minutes whereas feed D₁ had the lowest water stability of 18.54 ± 2.10 minutes. Feed D₂ had water stability of 32.76 ± 1.05 minutes, D₃ (35.23 ± 2.42 minutes) and D₄ (39.12±2.94).

Table 1. Percentage composition of experimental diets

Ingredients	Experimental diets				
	D ₁	D ₂	D ₃	D ₄	D ₅
Wheat bran	55.36	55.36	55.36	55.36	55.36
Fish meal	21.67	21.67	21.67	21.67	21.67
Soya bean	21.67	21.67	21.67	21.67	21.67
Premix	0.30	0.30	0.30	0.30	0.30
Vitamin c	0.05	0.05	0.05	0.05	0.05
Salt	0.30	0.30	0.30	0.30	0.30
Methionine	0.35	0.35	0.35	0.35	0.35
Lysine	0.30	0.30	0.30	0.30	0.30
Baobab leave	0	4	8	12	16

Table 2. Proximate composition of the experimental diets

Indices	D ₁ (0% BLM)	d ₂ (4% BLM)	d ₃ (8% BLM)	d ₄ (12% BLM)	d ₅ (16% BLM)
Crude protein	29.47±0.55 ^b	31.7±0.32 ^{ab}	32.05±1.05 ^a	31.44±0.45 ^{ab}	32.80±1.11 ^a
Fat	6.55±7.66 ^c	7.52±1.00 ^b	7.77±0.57 ^b	8.01±2.48 ^a	8.10±2.50 ^a
Fibre	13.33±2.51 ^c	16.00±1.0 ^{bc}	16.00±1.0 ^{bc}	17.00±2.30 ^{ab}	9.66±1.15 ^a
Ash	2.00±0.00 ^a	2.33±0.57 ^a	1.66±0.57 ^a	2.00±0.00 ^a	2.33±0.57 ^a
Dry matter	89.43±7.66 ^a	97.23±0.64 ^a	97.30±0.43 ^a	94.31±1.59 ^a	95.80±2.95 ^a
Moisture	10.56±7.66 ^a	2.76±0.64 ^a	2.70±0.43 ^a	5.60±1.51 ^a	4.28±0.95 ^a
Nfe	38.09±0.043	39.62±0.094	39.8±0.193	35.95±0.225	32.83±0.316

*means with the same superscript are not significantly different ($p>0.05$)

Table 3. Pellets characteristics of experimental diets formulated with baobab leaf meal (BLM)

Pellet characteristics	D ₁ (0% BLM)	D ₂ (4% BLM)	D ₃ (8% BLM)	D ₄ (12% BLM)	D ₅ (16% BLM)
Initial weight of pellets (g)	1.42±0.08 ^c	2.69±0.06 ^b	3.12±0.20 ^a	3.01±0.80 ^{ab}	3.40±0.15 ^a
Weight of retained whole pellets (g)	0.53±0.08 ^d	1.76±0.08 ^c	2.20±0.20 ^{bc}	2.36±0.23 ^b	2.90±0.05 ^a
Stability (mins)	18.54±2.10 ^c	32.76±1.05 ^b	35.23±2.42 ^b	39.12±2.94 ^{ab}	42.66±1.17 ^a
Floatability (mins)	0.00±0.00 ^d	8.33±2.88 ^c	10.00±5.00 ^c	25.00±0.00 ^b	41.66±2.88 ^a
Floatability rate (%)	0.00±0.00 ^d	16.66±2.88 ^c	20.00±5.00 ^c	50±0.00 ^b	83.32±2.88 ^a

*means with the same superscript are not significantly different ($p>0.05$)

Findings of this study showed that feed formulated with baobab leaf meal (BLM) exhibited floating ability which increased with the increase in inclusion level. The control feed (D₁ – 0% BLM) formulated with no baobab leaf meal (BLM) inclusion had no floating ability while D₅ (16% BLM) had the highest floating ability and water stability. In fish feed formulation, water stability, floatability and nutrient leaching rate are the main issues. Although the feed will sink and disintegrate but it is lower compared to the time taken for the fishes to consume the feed that is disintegrate but it is lower compared to the time taken for the fishes to consume the feed that is 10-15 minutes [1]. The implication of findings obtained in this study is that feeding fish with feed d₅ (16% BLM) will not result in loss of feed

pellet and nutrients due to sinking into mud or pond bottom which may decay leading to water pollution and bacterial growth which may cause diseases. The different inclusion level of baobab (*A. Digitata*) leaf added to the feed, contributed to the floatability and the stability of the fish feed after exposure for 50 minutes. According to Solomon et al. [16], wheat grain starch (WGS) recorded 50% floatation at 50 minutes exposure to water. This is however lower than results obtained for feed D₅ (16% BLM) with a floatability rate of 83.32% but similar to floatability rate of 50% obtained for feed D₄ (8% BLM) after exposure to water for 50 minutes. The difference could be attributed to the difference in the ingredients used in formulating the experimental diets. This implies that the inclusion of baobab

leaf meal (BLM) in fish feed will result in a better floatability compared to wheat grain starch (WGS). The floatability characteristics observed in baobab leaf meal (BLM) could be due to the presence of high gluten protein in baobab leaf meal (BLM) compared to wheat grain starch (WGS). Orire et al. [17] reported a water stability as high as 82.81% in fish feed formulated with cassava starch as a binder after 50 minutes exposure to water. This is however lower than the 83.32% reported for feed D₅ (16% BLM) in the present study. Obi et al. [18] reported a floatation period of 40% when crushed water melon shell was added at 15% in a fish diet, this is however lower than 50% obtained from this study at 12% inclusion of (BML). Findings of this study indicates that baobab leaf meal (BLM) has proven to aid feed buoyancy and stability when included in the right form and percentage. When feed sinks, there is a serious nutrient loss due to leaching of the essential vitamins like vitamin A, D, E, K of fat soluble status and about one third of the free plus protein bound amino acid. Extruded floating feed cost is quite a disadvantage over a dried and moist pellet [19]. And as such, floating feed is a management tool as it enables the farmer to observe the feeding activity of their fishes [20]. Though feed (D₁ – 0% BLM) and (D₂ – 4% BLM) exhibited low buoyancy, the two feeds can still be utilized by benthic feeders like catfish [21].

The result from this study showed that ingredients used in fish feed formulation influenced the pellet characteristics. The natural binding quality of the ingredient used in feed formulation could be utilized to their fullest capacity instead of adding non-nutritive agents. Therefore, to formulate floating local feed, careful selection of feedstuff or ingredients is a necessity to enhance the buoyancy of feed since some feedstuffs have positive buoyancy characteristics.

4. CONCLUSION

The use of baobab leaf meal (BLM) as a binder and floatability agent in local feed formulation has yielded a very positive result in the present study. Baobab leaf meal (BLM) is relatively cheap, toxic free and available specifically in the northern part of Nigeria. Baobab leaf meal (BLM) is easy to process and its usage in floating feed formulation is cheap compared to the cost of importing extruded floating feed from the western nation. However, there is a need to perform an in-vitro experiment with fish.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Masser MP, Wurts WA. Managing recreational fish ponds. World Aquaculture. 1992;23(2):41-47.
2. Lim C, Cuzon G. Water stability of shrimp pellet: A review. Asian Fisheries Science. 1994;7:115–127.
3. Ballarin AM, Heller Lo. Evaluation of yam starch (*Discorea rotundata*) as aquatic feed binder. Pakistan Journal of Nutrition. 2010;9(7):668–671.
4. Johnson TA, Wandsvick SK. Fish nutrition and development in aquaculture. Published by C & H. 2-6 Boundary Row, London SE1 8HN. 1991;1-119.
5. Sadiku, Jauncey K. Digestibility, apparent amino acid available and waste generation potential of soybean flour – poultry meat blend based diets for tilapia fingerlings. Aquaculture Research. Survey NIFFR, New Bussa, Nigeria Techniques on the Anti-nutrient Contents of Baobab Seeds (*adansonia digitata*). 1995;26:651–657.
6. Wood J. Selecting equipment for producing farm made aquafeed. In: New MB, Tacon Ajg, Savas C (eds) Farm Made Aquafeed. FAO/AADCP Thailand. 1993; 135-147.
7. Mgbenka BO, Lovell RT. The progressive fish culturist. Aquaculture Fish Service with U.S. Dept, IF & W. Series. 1984;46:4.
8. Gebauer J, El-Siddig K, Ebert G. Baobab (*Adansonia digitata* L.): A review on a multipurpose tree with promising future in the Sudan. 2002;67:155-160.
9. Huges AO, Phillips OM. Flow and reactions in permeable rocks. Cambridge Univ. Press, New York. 1989;285.
10. Yazzie D, Vanderjagt DJ, Pastuszyn A, Okolo A, Glew RH. The amino acid and mineral content of baobab (*Adansonia digitata* L.) leaves. J Food Compost Anal. 1994;7:189-193.
11. Sidibe M, Williams JT. Baobab, *Adansonia digitata*. Southampton: International Centre for Underutilised Crops; 1998.
12. Baum DA. A systematic revision of *Adansonia* (bombacaceae). Ann MO Bot Gard. 1995;82:440-470.

13. Baum DA. The ecology and conservation of the baobabs of Madagascar. Primate Rep. 1996;46:311-327.
14. Eyo VO, Ekanem AP, Jimmy UU. A comparative study of the Gonado-Somatic Index (GSI) and gonad gross morphology of African catfish (*Clarias gariepinus*) fed unical aqua feed and coppens commercial feed. Croatian Journal of Fisheries. 2014;72:63–69.
15. AOAC. Official methods of analysis. 15 Ed. The Association of Official Analytical Chemists, Washington, DC; 1990.
16. Solomon SG, Ataguba GA, Abeje A. Water stability and floatation test of fish pellets using local starch sources and yeast (*Saccharomyces cerevisiae*). International Journal of Latest Trends in Agriculture and Food Sciences. 2011;1(1).
17. Orire AM, Sadiku SO, Tiamioyu LO. Suitability of cassava starch as feed binder. Science Forum: J. Pure Appl. Sci. 2001;61-65.
18. Obi MN, Kolo RJ, Orire AM. The production of floating fish feed using melon shell as a floating agent. International Journal of Science and Nature. 2011;2(3): 477-482
19. Effiong BN, Sanni A, Sogbesan OA. Comparative studies on the binding potential and water stability of duckweed meal, corn starch and cassava starch. New York Science Journal. 1992;2(4):50–57.
20. Rokey G, Plattner G. A practical approach to aquafeed extrusion. Feed Mgmt. 2003;54(1):24–27.
21. Falayi BA. Comparative studies of binding agents for water stability and nutrient retention in African catfish *Clarias gariepinus* (B) feeds. Masters in Technology Thesis Dept. of Wildlife and Fisheries, Federal University of Technolgy, Akure. 2000;75.

© 2018 Felix and Oscar; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/26403>